

SULFATION-INDEPENDENT L-SELECTIN LIGAND  
(SILL) AND THERAPEUTICS THEREOF

Ind 9.1 >

This application is a continuation-in-part  
5 application of United States Serial Number 08/321,400  
filed October 11, 1994.

BACKGROUND OF THE INVENTION

10

TECHNICAL FIELD

15

The present invention involves the  
development of compounds which can regulate and  
control the function of adhesion molecules as well as  
methods and apparatus for testing for adhesion  
molecules.

BACKGROUND ART

20

25

Adhesion molecules are involved in the  
fundamental control of cell-cell interaction and  
cellular migration. Adhesion molecules regulate  
diverse processes in inflammation, hematopoiesis and  
tumor metastasis. (Woodruff, et al, 1987; Springer,  
et al, 1987; Sharon and Lis, 1993; Sackstein, 1993)  
For general reviews on adhesion molecules see Carlos  
and Harlan, 1994 and Chin et al, 1991. It would be

093316-072499

useful to develop reagents which can control and regulate the adhesion proteins, particularly within the selectin family.

The peripheral lymph node "homing receptor",  
5 L-selectin (CD62L), is a ~75 kDa glycoprotein which mediates attachment of lymphocytes to lymph node (LN) high endothelial venules (HEV), an adhesive interaction which is the first step in the migration of lymphocytes from blood into lymphoid tissues  
10 (Gowans and Knight, 1964; Marchesi and Gowans, 1964).

This trafficking of lymphocytes from blood into lymph nodes is markedly nonrandom and is initiated by specific adherence of the lymphocytes to HEV. L-selectin is the principal lymphocyte membrane  
15 glycoprotein mediating this attachment (see Shimizu and Shaw, 1993 and also United States Patent 5,489,578 columns 3-4 for a review).

The L-selectin protein is recognized by a variety of monoclonal antibodies (mAbs) in humans  
20 (Gatenby et al., 1982 (Leu-8); Reinherz et al., 1982 (TQ-1); Tedder et al., 1990 (LAM)) and is a member of the selectin family of adhesion molecules, which includes P-selectin (CD62P) and E-selectin (CD62E). Selectins share a common structure consisting of an  
25 amino-terminal calcium-dependent lectin domain, an epidermal growth factor domain, a variable number of repeat sequences bearing homology to complement

regulatory and catalytic proteins binding C3b or C4b, a transmembrane portion, and a C-terminal cytoplasmic tail (Bevilacqua and Nelson, 1993; Rosen, 1993). The molecular weight varies among leukocytes due to differences in posttranslational glycosylation among subsets of leukocytes (Carlos and Harlan, 1994). The lectin domain of these proteins directs their adhesion to carbohydrate molecules present on the cell surface.

The adhesive interaction between lymphocytes and HEV has been extensively analyzed using an *in vitro* binding assay (Stamper and Woodruff, 1976). The adhesive interaction between L-selectin and its ligand(s) was first identified by this *in vitro* binding assay in which lymphocytes are overlaid onto frozen sections of lymph nodes (Stamper and Woodruff, 1976). This "lymphocyte-HEV adherence assay" is an *in vitro* approximation of physiologic adhesion mediated by L-selectin and has been the conventional approach for studying the adhesive function of L-selectin in its native state on the surface of leukocytes. Indeed, in early studies, antibodies directed against L-selectin were operationally identified by their ability to block lymphocyte-HEV adherence in the assay (Rasmussen et al., 1985; Gallatin et al., 1983), and these antibodies subsequently led to the identification of L-selectin at a molecular level. This assay is performed under shear at 4°C, whereby

binding mediated by L-selectin is maximized and effects of other adhesion molecules are minimized (Shaw et al, 1986; Spertini et al., 1991). Under these conditions, lymphocytes will adhere specifically to lymph node HEV via interactions between L-selectin and its corresponding ligand(s).

The interaction of L-selectin with its corresponding ligand(s) on HEV is calcium-dependent (Woodruff et al., 1977) and requires the presence of sialic acid (Rosen et al., 1985; True et al., 1990) and sulfate (Imai et al., 1993) on the ligand(s) (see United states Patent 5,489,578 column 4 for a review).

L-selectin behaves as a lectin and recognizes sialylated, high mannose residues on its corresponding endothelial ligands which are identified by the monoclonal antibody MECA-79 (Streeter, 1988; ATCC Accession number HB-9479; United States Patent 5,403,919).

Most of what is known about L-selectin ligands has been derived from studies of molecules recognized by MECA-79 (Streeter et al, 1988; ATCC Accession Number HB-9479; United States Patent 5,403,919), a rat anti-murine lymph node monoclonal antibody which identifies HEV by attaching to L-selectin ligands. In murine lymph nodes, MECA-79 and a murine L-selectin-Ig chimeric construct each immunoprecipitate several glycoproteins (Watson et

al., 1990), two of which have been well-characterized: GLYCAM-1 (Imai et al., 1991), a secreted protein of 50,000 m.w., and CD34, an integral membrane protein of ~90,000 m.w. (Baumhueter et al., 1993). *In vitro* adherence of lymphocytes via L-selectin can be inhibited by carbohydrates such as mannose-6-phosphate (man-6-P), PPME (Phosphomannan monoester core from *Hansenula hostii*, a phosphomannosyl-rich polysaccharide), and fucoidin (a sulfated, fucose-rich polysaccharide) (Stoolman and Rosen, 1983). In addition, the MECA-79 determinant is found on an HEV glycoprotein of 200,000 m.w. (Hemmerich et al., 1994) and on a selectively modified subset of the mucosal vascular addressin (MAdCAM-1) which supports L-selectin-mediated lymphocyte adhesion under shear (Berg et al., 1993). The HEV ligands recognized by MECA-79 are all heavily glycosylated mucin-like proteins (Shimizu and Shaw, 1993) rich in O-linked glycosylations bearing sialic acid. Treatment of endothelial L-selectin ligands with neuraminidase (Hemmerich et al., 1994; True et al., 1990) and with O-sialoglycoprotease abrogates the binding of L-selectin (Puri et al., 1995).

The molecular determinant recognized by both MECA-79 and L-selectin chimera is sulfation-dependent as demonstrated by markedly diminished ligand recognition following metabolic inhibition of

5 sulfation by chlorate (Hemmerich et al., 1994). These results have highlighted the critical role of sulfation in ligand activity (Hemmerich et al., 1994; Imai et al., 1993), a conclusion supported by the fact that naturally-occurring unsulfated GLYCAM-1 is not capable of binding L-selectin (Dowbenko et al., 1993).

10 Moreover, sulfation is characteristic of purified glycolipids (Suzuki et al., 1993), oligosaccharides (Greene et al., 1995) and glycosaminoglycans (Stoolman and Rosen, 1983) which bind to L-selectin, and the degree of sulfation correlates with binding affinity (Stoolman and Rosen, 1983).

15 Although present on endothelial cells in most tissues (Beschorner et al., 1985), CD34 is best known for its expression on the earliest multilineage colony-forming hematopoietic stem cells (Civin et al., 1984). Hematopoietic progenitor cells characteristically express both L-selectin and CD34 (Terstappen et al., 1992), and there is growing  
20 evidence that L-selectin plays a role in hematopoiesis (Terstappen et al., 1993; Kobayashi et al., 1994; Dercksen et al., 1995). The characterization of L-selectin and its ligands among progenitor cells is of considerable interest as adhesion proteins regulate  
25 cell-cell and cell-stromal interactions fundamental to hematopoiesis.

It would be useful to have strategies which would allow regulation of hematopoiesis since it is regulated by cell-cell and cell-stromal interactions.

For example, Terstappen et al (1993) have shown that  
5 activation of L-selectin increases the clonogenic  
capacity of stem cells. Moreover, Dercksen et al  
(1995) have demonstrated that hematologic recovery  
following hematopoietic stem cell transplantation in  
humans correlates with the number of CD34+/L-selectin+  
10 cells infused.

During recovery of immune function following  
bone marrow transplantation, pathologic changes have  
been observed following transplantation which  
interfere with lymphocyte migration and HEV integrity.

15 Further, in addition to changes in lymph node  
structure, alterations in lymphocyte migration can  
occur secondary to the effect of pharmacologic agents  
used in posttransplant therapy such as corticosteroids  
(Sackstein, 1993). It would be useful to have an  
20 agent which can assist in reestablishing lymphocyte  
trafficking and so immune function following bone  
marrow transplantation.

The crucial role of adhesion molecules in  
controlling and directing the inflammatory process  
25 indicates that a reagent which interferes with the  
process, i.e., an anti-adhesive (see for example  
United States Patents 5,489,578; 5,512,442;

036116-07249  
66720-913650

5,304,640), could have anti-inflammatory properties.

Further, cell adhesion molecules are involved in metastasis, therefore it would be useful to develop an anti-adhesive which has anti-metastatic properties. In particular, with the identification of L-selectin on hematopoietic cells, it would be useful to have an anti-adhesive that affects L-selectin in leukemia to decrease the growth and spread of malignant hematopoietic cells throughout the body.

Further, it would be useful to have additional cell markers and monoclonal antibodies directed against these cell markers to allow for cell targeting.

In general, assays for determining the adhesion between cells based on shear require the use of frozen sections of the substrate or target cell (Stamper and Woodruff, 1976; Sackstein et al, 1988) as described hereinabove and a cell suspension overlaying the substrate. This conventional overlay adherence/binding assay has previously been used only with frozen tissue sections, with lymph nodes as a principal example, as the substrate. The frozen section has the advantage of being a predictable monolayer. A monolayer is required since the overlay adherence assay mimics the blood flow over a vessel wall.



It would be useful to have available isolated cells or cell lines as the substrate in the assay. This would enable the use of cell suspensions rather than tissue blocks, giving rise to more reproducible results as well as increasing the types of cells available as a target and reducing the need for surgical procedures for tissue removal.

Initially, applicants attempted to grow the cell lines directly on the glass slides. However, the assay did not produce reproducible results because a predictable, consistent monolayer of cells was not generated. If there is no consistent monolayer, i.e. if there is an uneven surface due to clumping etc. of the cells, then adherence artifacts are found.

Initial attempts to use preparations of single cell suspensions deposited on a glass slide utilizing a Cytospin 3 Cytocentrifuge (Shandon Lipshaw, Pittsburgh, PA) were also generally unsuccessful. Procedures using a cytocentrifuge placed a single layer cell pellet/substrate in the center of the slide (Figure 11, 13, 15). When these slides were used in the assay on a rotating platform (e.g. 80 rpm rotation) in the adherence assay at 4°C for 30 minutes the rotational shear forces placed the overlay cell suspension utilized (typically lymphocytes) at a variety of positions on the slide depending on the platform itself, the platform angle

09358115-022199

and the rotation speed as well as other factors. In general, the overlay cell suspension was not localized over the center of the slide above the cytocentrifuged placed cell substrate. It would therefore be useful to be able to modify the placement of the cytocentrifuged cell pellet (to any selected location on the slide) so that it can be used as a substrate in an adherence assay dependent on the shear conditions employed for any given assay.

#### SUMMARY OF THE INVENTION

According to the present invention, an isolated and purified glycoprotein and functional analogues thereof are disclosed. The glycoproteins are characterized by being expressed on at least primitive hematopoietic cells, and being a ligand for L-selectin. The binding of ligand to L-selectin is sulfation-independent and it is neither inhibited by anti-CD34 antibodies nor by MECA-79 monoclonal antibody and is resistant to O-sialoglycoprotein endopeptidase activity.

Further, the present invention provides a method of performing an overlay adherence assay by using isolated cells or cell lines as a substrate. The cells are prepared as the substrate for the assay using a cytocentrifuge with a modified sample chamber

allowing placement of the cytocentrifuged cell pellet to any selected location on the slide determined by the shear conditions employed for any given assay.

5

#### BRIEF DESCRIPTION OF THE DRAWINGS

Other advantages of the present invention will be readily appreciated as the same becomes better understood by reference to the following detailed description when considered in connection with the accompanying drawings wherein:

FIGURE 1A-B are photomicrographs of cytocentrifuge preparations of KG1a cells demonstrating adherence of lymphocytes (small dark dots), (A) Lymphocytes adhere to KG1a in the presence of CD45 or isotype control antibodies (Abs), (B) Lymphocyte binding assay in the presence of LAM1-3 antibody (anti-L-selectin);

FIGURE 2A-C are FACS profiles of lymphocytes used in the binding assay after incubation with (A) isotope-matched IgG control, (B) LAM1-3, or (C) anti-CD45 antibodies, followed by GAM-FITC, results shown are representative of three independent experiments;

FIGURE 3A-D are FACS histogram profiles of KG1a cells sorted by FACS prior to the binding assay into CD34+ (A,B) and CD34- (C,D) fractions using mAb HPCA-2PE, control sorted cell fraction (A,C) were

09358146-072199

restained with a FITC-conjugated isotype matched monoclonal antibody, experimental sorted cell fractions were restained for CD34 using mAb QBEND10-FITC with (B) positive (>90%) and (D) negative (<10%) for CD34, results shown are representative of three independent experiments;

FIGURE 4 is a photomicrograph showing the results of lymphocyte adherence assay performed on the sorted cells, and no differences in lymphocyte adherence were evident among the CD34+ and CD34- populations, adherence to the CD34 negative fraction is shown;

FIGURE 5A-F are FACS profiles of COS-7 cells transfected with either CD34-pCDM8 (E,F) or pCDM8 (mock, C,D), then analyzed by FACS and compared to KG1a (A,B) for CD34 expression, antibodies used were isotype-matched IgG<sub>1</sub> control (ACE) and anti-CD34 mAb QBEND10 (BDF), lymphocytes did not adhere to CD34-transfected COS-7 cells, despite higher levels of CD34 expression as compared to KG1a cells;

FIGURE 6A-C wherein: (A-B) are FACS profiles of KG1a cells stained with IgM control antibody (A, top) or MECA-79 (B, bottom) followed by fluorescein isothiocyanate (FITC)-conjugated goat anti-rat IgM and Figure 6 (C) is a photograph of a gel showing KG1a and murine lymph node (LN) metabolically radiolabeled with <sup>35</sup>SNa<sub>2</sub>SO<sub>4</sub> and immunoprecipitated with MECA-79, murine

anti-human CD43 as a positive control or respective isotype control antibodies;

FIGURE 7A-C are photographs of representative results of adherence assays using suspensions of rat thoracic duct lymphocytes overlaid onto cytocentrifuge preparations of KG1a cells of light microscopy (250x) of lymphocytes (dark circles) adhering to cytocentrifuge preparations of KG1a cells following incubations with RPMI containing buffer alone (A, top), immediately following neuraminidase treatment (B, middle), and neuraminidase treatment followed by twenty-four hour incubation in 10 mM sodium chlorate (C, bottom), note the absence of lymphocyte adherence immediately following neuraminidase treatment;

FIGURE 8A-C are graphs showing representative results of flow cytometric analysis of sialic acid-dependent epitope of CD43 demonstrating recovery of surface CD43 despite inhibition of sulfation by indirect immunofluorescence, wherein (A) KG1a control levels, (B) KG1a immediately following neuraminidase treatment, and (C) KG1a 24 hours following neuraminidase treatment, cells were cultured in the presence of 10 mM sodium chlorate, monoclonal antibody L60 was used as primary antibody and the secondary antibody was PE-conjugated;

FIGURE 9 A,B are photographs of gels wherein (A) shows the inhibition of sulfation by sodium chlorate demonstrated by immunoprecipitation/SDS PAGE analysis of  $^{35}\text{S}$ - $\text{SO}_4$  incorporation into CD43 and (B) is an autoradiograph of SDS-PAGE analysis of total  $^{35}\text{S}$ - $\text{SO}_4$ -radiolabelled and  $^{35}\text{S}$ -methionine/cysteine-d protein in the presence (+) and absence (-) of chlorate, lanes contain equivalent amounts of lysate material obtained from identical numbers of cells cultured in the presence of respective radiolabels for the terminal eight hours of the 24 hour culture period, marked inhibition of sulfation is demonstrated among all sulfated proteins in the presence of chlorate, without significant effects on protein synthesis as shown by equivalent profiles of  $^{35}\text{S}$ -methionine/cysteine-radiolabelled proteins, molecular weight markers in kDa are shown at the left of the Figure;

FIGURE 10 is an exploded view of a holder assembly showing a modified sample chamber of the present invention;

FIGURE 11 is a front view of an assembled holder assembly of the prior art;

FIGURE 12 is a front view of an assembled holder assembly of the present invention showing the lateral and vertical offset of the modified sample chamber;

FIGURE 13 is a top plan view of the assembled holder assembly as shown in FIG. 11;

FIGURE 14 is a top plan view of the assembled holder assembly as shown in FIG. 12;

5           FIGURE 15 is a side view of the assembled holder assembly as shown in FIG. 11;

FIGURE 16 is a side view of the assembled holder assembly as shown in FIG. 12;

10           FIGURE 17 is a front view of a slide marked for use in the present invention; and

15           FIGURE 18 is a front view of a holder assembly with a sample chamber of the present invention partially cut away to show cellular deposition at the position indicated on the slide in FIG. 17.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

20           The present invention provides an isolated and purified glycoprotein that is a sulfation-independent L-selectin ligand (SILL) and functional analogues thereof.

25           The term analogue as used herein is defined as a SILL variant (alternatively the terms alteration, amino acid sequence alteration, amino acid sequence variant, glycosylation variant can be used) with some differences in their amino acid sequences or

095815-0419  
66T220-9T3560

glycosylation pattern as compared to the native sequence of the SILL. Ordinarily, an analogue will be generally at least 70% homologous with the native SILL over any portion that is functionally relevant. In more preferred embodiments the homology will be at least 80% and can approach 95% homology to the amino acid sequence of the protein segment of SILL. The homology will extend over a region of contiguous amino acids. The amino acid sequence of an analogue may differ from that of the glycoprotein of the present invention when at least one residue is deleted, inserted or substituted.

Differences in glycosylation may be present between the analogue and the present invention (see United States Patent 5,489,578 column 1, lines 34-67 for a review of carbohydrates in physiologically relevant recognition/adhesion). For glycoproteins, the post-translational sugar modification are, in some cases, the functional determinants of the molecule. The core protein sequence may be important in the spatial display of the sugars but may not contribute to the activity directly. Ordinarily, an analogue will be generally at least 70% homologous with the native SILL over any portion that is functionally relevant. In more preferred embodiments the homology will be at least 80% and can approach 95% homology to the glycosylation pattern of the carbohydrate portion



of SILL. The molecular weight of the glycoprotein may vary between the analogue and the present invention due to carbohydrate differences.

Functionally relevant refers to the

5 biological property of the molecule and in this context means an *in vivo* effector or antigenic function or activity that is directly or indirectly performed by a naturally occurring (native) SILL. Effector functions include but are not limited to

10 include receptor binding, any enzymatic activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to extracellular matrix or cell surface molecules, or any

15 structural role. The antigenic functions essentially mean the possession of an epitope or antigenic site that is capable of cross-reacting with antibodies raised against a naturally occurring SILL.

Biologically active SILL analogues share an effector

20 function of the native SILL which may, but need not, in addition possess an antigenic function.

The glycoprotein SILL has the following functional characteristics. It is expressed on at least primitive hematopoietic cells. The glycoprotein

25 is a ligand for L-selectin. The ligand binding to L-selectin is not inhibited by anti-CD34 antibodies and is not recognized by the MECA-79 monoclonal antibody.

00358446-072199

The glycoprotein exhibits sulfation-independent function. The glycoprotein maintains L-selectin ligand function following treatment with O-sialoglycoprotein endopeptidase. The glycoprotein is  
5 designated hereinafter as sulfation-independent L-selectin ligand, SILL.

Further, SILL, is a membrane associated glycoprotein and functions as an adhesion protein ligand. The glycoprotein mediates attachment of L-  
10 selectin expressing cells such as lymphocytes to hematopoietic cells including primitive hematopoietic cells.

SILL was initially identified on KG1a and is similar to endothelial L-selectin ligands in that its  
15 activity is calcium-dependent, requires the presence of sialic acid for L-selectin binding, and it is a glycoprotein. However, SILL differs in two fundamental ways from its endothelial counterparts as currently characterized, and represents a novel L-  
20 selectin ligand glycoprotein. Firstly, although both mouse and human lymphocytes bind to KG1a through an L-selectin-mediated interaction (Rosen, 1994), neither a murine L-selectin-Ig chimera (LEC-IgG) nor a human L-selectin-IgG chimera attach to KG1a cells, though  
25 these molecules recognize endothelial L-selectin ligands (Majdic, 1994).

Secondly, the KG1a L-selectin ligand, SILL, does not contain MECA-79-recognized epitopes. Both flow cytometric analysis of KG1a (SILL) and immunoprecipitation studies of radiolabeled KG1a membrane proteins show no evidence of MECA-79 antigens on the surface of KG1a cells (Figures 6A, 6B), indicating that this epitope is not involved in the binding determinants of the KG1a L-selectin ligand (SILL). Enzymatic cleavage studies and membrane stripping studies under a variety of pH and salt conditions indicate that the L-selectin ligand expressed on KG1a is a non-GPI-linked integral membrane glycoprotein.

Applicant asked whether the ligand of the present invention has the central structural features of other naturally-expressed L-selectin ligands, all of which have been characterized on endothelial cells.

The role of sulfation in the binding activity of the KG1a L-selection ligand is at the core of this question, since it was shown in two previous "landmark" reports (Imai et al., 1993; Hemmerich et al., 1994) that the function of endothelial L-selectin ligands is sulfation-dependent. The Hemmerich et al. article in particular describing an L-selectin ligand, emphasized the functional role of sulfation for all endothelial L-selectin ligands, whereas the Imai et al. article focused only on the sulfation-dependent

function of the secreted endothelial L-selectin ligand known as GlyCAM-1. Applicant's new data clearly distinguishes the L-selectin ligand of the present invention in two important ways: (1) it is not  
5 recognized by monoclonal antibody MECA-79, an antibody which recognizes the binding domain of all previously identified membrane L-selectin ligands, and (2) it does not require sulfation for binding to L-selectin.

Thus, Applicant has unequivocally established that  
10 this ligand is novel - not only in the fact that it is expressed on a hematopoietic cell line but that it is structurally unique from all previously identified endothelial ligands.

Further, there is an independent  
15 relationship between P-selectin glycoprotein ligand-1 (PSGL-1) and the SILL ligand activity showing that they are distinct molecules (Example 7).

The present invention also provides for an antibody directed against SILL. The antibodies may be  
20 either monoclonal or polyclonal. Murine monoclonal antibodies are initially raised against KG1a cells. The monoclonals that are generated are then screened for the ability to block lymphocyte binding to KG1a. (See also United States Patent 5,130,144 which  
25 discloses a method in which a monoclonal antibody is raised against KG1a cells.)

0935016.0219

Utilizing these monoclonal antibodies, the SILL is isolated by immunoprecipitation of KG1a membrane lysates as is standard in the art and used for the production of further antibodies as needed.

5 Such methods can be found described Sambrook et al, *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor, New York, 1989, as well as additional methods of isolation and purification as are known in the art.

667229 978560  
0935315 072199

10 Additionally, the antibodies may be prepared against a synthetic peptide based on the sequence, or prepared recombinantly by cloning techniques or the natural gene product and/or portions thereof may be isolated and used as the immunogen. Such proteins or peptides can be used to produce antibodies by standard  
15 antibody production technology well known to those skilled in the art as described generally in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988 and Borrebaeck, *Antibody Engineering - A Practical Guide*,  
20 W.H. Freeman and Co., 1992. Antibody fragments may also be prepared from the antibodies and include Fab, F(ab')<sub>2</sub>, and Fv by methods known to those skilled in the art.

25 For producing polyclonal antibodies a host, such as a rabbit or goat, is immunized with the protein or peptide fragment of the protein, generally with an adjuvant and, if necessary, coupled to a

carrier; antibodies to the protein are collected from the sera.

For producing monoclonal antibodies the technique involves hyperimmunization of an appropriate donor with the protein or peptide fragment, generally a mouse, and isolation of splenic antibody producing cells. These cells are fused to a cell having immortality, such as a myeloma cell, to provide a fused cell hybrid which has immortality and secretes the required antibody. The cells are then cultured, in bulk, and the monoclonal antibodies harvested from the culture media for use.

The antibody or antibody fragment can be bound to a solid support substrate or conjugated with a detectable moiety or be both bound and conjugated as is well known in the art. (For a general discussion of conjugation of fluorescent or enzymatic moieties see Johnstone & Thorpe, *Immunochemistry in Practice*, Blackwell Scientific Publications, Oxford, 1982.) The binding of antibodies to a solid support substrate is also well known in the art. (see for a general discussion Harlow & Lane *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Publications, New York, 1988) The detectable moieties contemplated with the present invention can include, but are not limited to, fluorescent, metallic, enzymatic and radioactive markers such as biotin, gold, magnetic

0935846-018560

particles, ferritin, alkaline phosphatase,  $\beta$ -galactosidase, peroxidase, urease, fluorescein, rhodamine, tritium,  $^{14}\text{C}$  and iodination.

5 The method of targeting cells includes the steps of preparing antibodies directed against SILL as described above and coupling the antibodies to the appropriate agent whether for cell killing, cell selection or cell identification. For cell killing, toxins such as ricin A chain, pseudomonas exotoxin A, 10 diphtheria toxin, other plant and bacterial toxins as well as chemotherapeutic compounds can be coupled in the present invention forming an immunotoxin. For a general review of the antibody-toxin art see Ramakrishnan, 1990.

15 Cell targeting requires exposing a population of cells to the immunotoxin. A toxin bound antibody can be administered to the appropriate patient and targeted cells killed *in vivo*. The immunotoxin is administered and dosed in accordance 20 with good medical practice, taking into account the clinical condition of the individual patient, the site and method of administration, scheduling of administration, and other factors known to medical practitioners. The "effective amount" for purposes 25 herein is thus determined by such considerations as are known in the art. The amount must be effective to achieve improvement including but not limited to

0035516-0220-9T55550

improved survival rate, more rapid recovery, or improvement or elimination of symptoms.

5 A pharmacological formulation of the antibody can be administered to the patient in an injectable formulation containing any compatible carrier, such as various vehicle, adjuvants, additives, and diluents; or the compounds utilized in the present invention can be administered parenterally to the patient in the form of slow-release  
10 subcutaneous implants of the targeted monoclonal antibodies, or iontophoretic, polymer matrices, liposomes, and microspheres can be employed. Examples of delivery systems useful in the present invention include: U.S. Patent Numbers 5,225,182; 5,169,383;  
15 5,167,616; 4,959,217; 4,925,678; 4,487,603; 4,486,194; 4,447,233; 4,447,224; 4,439,196; and 4,475,196. Many other such implants, delivery systems, and modules are well known to those skilled in the art.

Alternatively, cells can be removed from the  
20 patient and treated ex vivo selectively. For example, cells expressing SILL can be removed through complement-mediated lysis from the ex vivo population and the remaining cells returned to the patient. Additional cell removal can be undertaken utilizing  
25 cell sorting, "panning" and magnetic bead separation.

Alternatively, utilizing cell sorting, "panning", magnetic bead separation and the like cell populations



can be enriched for SILL bearing cells and this enriched cell population returned to the patient if needed.

5 The targeted cells to be removed are cells expressing SILL and can be selected from the group consisting of leukemic cells, malignant hemopoietic progenitor cells, or any malignant cell expressing the marker.

667270 9118550  
10 The present invention also provides a method of regulating hematopoiesis, particularly in reconstitution of the immune system following bone marrow transplantation. The present invention includes the steps of selecting those cells with high(+) or low(-) expression of SILL depending on the  
15 growth characteristics associated with the marker density needed by the patient. The selection procedure utilizes ex vivo methods as described herein. After selection, the selected cell type is cultured in vitro, if needed, to expand the population  
20 using standard methods known in the art. The patient is then infused with the expanded, enriched SILL+ or SILL- population as needed.

25 The present invention further provides a method of regulating inflammatory response by interrupting cellular migration into lymph nodes and sites of both acute and chronic inflammation including the step of administering to the patient either

functional analogues or antibody directed against SILL, thereby disturbing cellular migration mediated by SILL by blocking the L-selectin-mediated cell attachment site and can be injected directly at the inflamed site if needed. The regulation of the inflammatory response would be useful in autoimmune disorders, post-ischemic tissue injury and sepsis (Carlos and Harlan, 1994). Administration and effective dose are as described for immunotoxins hereinabove.

See also United State Patents 5,484,891; 5,489,578; 5,464,778; 5,304,640; 5,538,724 and 5,403,919 for further methods and discussion of treatments with selectin ligands, all incorporated by reference in their entirety.

As described herein the adhesive interaction between cells is analyzed using a modified *in vitro* binding assay (Stamper and Woodruff, 1976). The *in vitro* binding assay requires that a cellular suspension be overlaid onto frozen tissue sections (Stamper and Woodruff, 1976). The assay requires that the substrate be positioned such that the forces of the rotating platform will deposit the overlay cell suspension on top of the substrate.

The present invention provides for the use of isolated cells instead of frozen tissue sections as the substrate of the binding assay. The cells may be established cell lines or isolated cells from any

source, mature or immature, including peripheral blood samples, from bone marrow aspirates, and from tissues donated through organ harvests or other sources as appropriate.

5           To prepare the cell substrate a protocol is used as further described in Example 1, which requires a modification of an existing sample chamber for use in a cytocentrifuge. When the prior art sample chamber as shown in Figures 11, 13 and 15 is used the  
10 cell substrate pellet from the cytocentrifuge is deposited in the middle of the slide to be used as the substrate. As described herein above this is not the preferred location for an overlay adherence assay. The preferred location must be determined for each  
15 rotation table and the substrate deposited at that site.

United States Patents 4,391,710, 4,678,579 and 4,729,778 disclose the cytocentrifuge and holder assembly of the prior art. The design and structure  
20 of a cytocentrifuge and the general design and construction of the holder assembly and sample chamber are disclosed in these patents and are incorporated in their entirety by reference. The terms used in describing a preferred embodiment of a modified sample  
25 chamber of the present invention are the same as in these patents where possible.

5 The present invention allows the use of  
isolated cells or cell lines as a substrate, the cells  
being able to be deposited on a selected location of a  
slide which must be determined for the shear  
conditions employed for any given adherence assay as  
described herein above. The present invention  
provides a sample chamber assembly comprising a sample  
chamber including cell substrate receiving means for  
receiving a cell substrate and depositing means for  
10 placing the cell substrate on a slide surface during  
cytocentrifugation, connecting means for connecting  
together a slide and sample chamber and fixing the  
depositing means at one of a plurality of positions  
relative to the connecting means. The present  
15 invention further provides a method of making a sample  
chamber assembly by connecting together a slide and a  
sample chamber, the sample chamber including a cell  
substrate depositing port and fixing the cell  
substrate depositing port at one of a plurality of  
20 positions relative to the slide.

In an embodiment, a sample chamber,  
generally shown at 20, is shown in Figure 10 which is  
an exploded view of a sample chamber assembly,  
generally shown at 22. The sample chamber assembly 22  
25 includes a holding member generally shown at 24, a  
slide 26, a filter card 28 and the sample chamber 20.  
The holding member 24 is also referred to as a slide

0935816-0229

clip.

5 The sample chamber 20 generally comprises a flat, generally rectangular plate 30 having a rectangular end flange 32 disposed normal to and along an edge of the plate 30 such that the intersection of the rectangular plate divides the end flange into two side areas, 33,33'. The rectangular end flange 32 and plate 30 are reinforced and interconnected by a generally triangular upper end piece 34 as best seen in Figure 10. In the prior art as shown in Figures 11 and 15 there is a lower end piece 36 opposite and spaced from the upper end piece 34, which is optionally present in the present invention. A vertical directed funnel 38 is contained within the plate 30 and interconnects with a horizontally extending discharge port 40 which extends from the funnel 38 within the plate 30 to the end flange 32. The discharge port 40 terminates at an opening 42 in the end flange 32. In the prior art sample chamber the distance between the opening 42 of the discharge port 40 and each of opposite long sides 44, 44' of the rectangular end flange 32 is equidistant. In other words in the prior art the two side areas 33,33' are generally the same size.

25 As described in the 4,391,710, 4,678,579 and 4,729,778 patents cell suspensions are loaded in the funnel 38 and deposited on the surface of the slide 26

093516-0229  
66 FEB 20 9 18 56

opposite the opening 42 of the discharge port 40.

5 A filter card 28 is disposed between the end flange 32 and the slide 26. The filter card 28 is dimensioned to be coextensive with the end flange 32 and is provided with an opening 46 that is coextensive with the opening 42 of the discharge port 40. In an embodiment the filter card 28 may have two openings 42 positioned such that the filter card 28 may be disposed between the end flange 32 and slide 26 in  
10 either vertical orientation such that the filter card opening 46 will be coextensive with the opening 42 of the discharge port 40. During operation the filter card 28 will absorb the liquid content of the cell sample and the cells will be deposited on the slide  
15 26. The filter card 28 can be separate as shown in Figure 10 or can be formed as part of the sample chamber 20 as described in United States Patent 4,678,579.

In the present invention the end flange 32  
20 includes opposite lateral edge regions 48, 48'. The lateral edge regions 48, 48' are disposed along each of the long sides 44, 44' of the rectangular end flange 32. In the present invention one of the lateral edge regions 48 is removed as best shown in  
25 Figures 12, 14 and 18. The removal of one of the lateral edge regions 48 allows the lateral displacement or offset of the sample chamber 20 in the

0152.00344-30

sample chamber assembly 22 as best shown in Figure 12, 14 and 18. To allow removal of one of the lateral edge region 48 the lower end piece 36 may have to be removed. The lateral displacement is in the direction of the removed lateral edge region 48 removed. Further the amount of one of the lateral edge region 48 removed is generally such that the distance from the opening 42 of the discharge port 40 in the end flange 32 to the resulting long side 44 of the end flange 32 after removal of the lateral edge region 48 generally corresponds to A as shown in Figure 17 as will be described and is no longer equidistant to the opposite long side 44' of the rectangular end flange 32. In other words the two side areas 33, 33' are no longer the same size, and in this embodiment 33 is smaller than 33'. Alternatively, the lateral edge region 48' can be removed. The filter card 28 is modified to be co-extensive with the end flange 32 after one of the lateral edge regions 48, 48' is removed.

The holding member 24 includes a generally U-shaped clip 50 which pivots over the slide 26, filter card 28 and sample chamber 20 when they are positioned in the holding member 24 to hold them in position in the holding member 24. The front surface of the end flange 32 is provided with horizontally directed, slotted ledges 52 on opposite sides of the

discharge port 40. The clip 50 when locked into position to hold the sample chamber assembly 22 together is disposed on the ledges 52 in the prior art as best shown in Figures 11 and 15. In the present invention when the sample chamber 20 is moved laterally, the clip must be disposed on both sides of the discharge port 40 and generally on the ledges 52 as best shown in Figure 12.

In the present invention the end flange 32 and rectangular plate 30 include a coextensive lower edge region 54. The lower edge region 54 is disposed along the lower short side of the rectangular end flange 32 and disposed along the lower edge of the rectangular plate 30 and spaced and parallel to the discharge port 40. In an embodiment of the present invention the lower edge region 54 is removed as best shown in Figures 16 and 18. The removal of the lower edge region 54 allows the vertical displacement or offset of the sample chamber 20 in the sample chamber assembly 22 as best shown in Figure 16. Further the amount of the lower edge region 54 removed is generally such that the distance from the center of the discharge port opening 42 to the resulting lower edge after removal of one of the lower edge region 54 generally corresponds to A' as shown in Figure 17 as will be described.



In the present invention when the sample chamber 20 is displaced vertically, the clip 50 while disposed on both sides of the discharge port 40 will generally not be disposed on the ledges 52 as best shown in Figure 16 but directly on the front surface of the end flange 32.

In an embodiment one of a lateral edge region 48, 48' and a lower edge region 54 are removed allowing a combined vertical and lateral displacement of the sample chamber 20 as best shown for one embodiment in Figure 18.

The modified sample chamber 20 can be made either by adapting existing prior art sample chambers formed in a plastic or other material which can be cut or shaved to size or the modified sample chamber 20 can be formed utilizing the methods disclosed in United States Patents 4,391,710, 4,678,579 and 4,729,778 and methods known to those skilled in the art of plastic extrusion or forming in other materials. A series of modified sample chambers can be prepared having incremental changes in the amount of lateral edge regions 48, 48', which opposite lateral edge region 48, 48' is elected for modification and/or lower edge region 54 removed allowing for different A and A' distances and prepared as a kit. The incrementally modified chambers are then available for testing to determine which

005315-07239  
661220-9718560

incremental change will deposit the cell substrate at the proper position on the slide.

Alternatively, the holding member 24 can be adapted to allow displacement of the slide 26. The holding member 24 is comprised of a channel 58, which can be U-shaped or M-shaped, having opposite side plates 60, a common web 62 interconnecting the opposite side plates 60 and a pivot rod 66 for interconnecting to the centrifuge. A retaining hook 64 is disposed on each side plate 60 to retain the clip 50 in the locked position. To adapt the holding member 24, one of the side plates 60 can be bent outwardly such that the slide 26 can be displaced laterally while the sample chamber 20 is either not moved or is moved oppositely laterally and/or vertically. This allows for maximum freedom of positioning of the cell pellet 56 on the slide 26. In adapting the side plate 60, care must be taken to insure that the retaining hook 64 can still be used.

The depositing means include the discharge port 40 and the discharge port opening 42 through which the cell substrate pellet 46 is deposited on the slide 26 at one of a plurality of positions. The depositing means can further include the filter card 28 and filter card opening 46 which is coextensive with the discharge port opening 42.

035515-07199  
667270-9773560

The receiving means include the funnel 38 interconnected with the discharge port 40 such that under cytocentrifugation a cell substrate 56 disposed in the funnel 38 enters the discharge port 40.

5           The connecting means comprise the holding member 24, end flange 32 and rectangular plate 30 as well as the lateral edge regions 48,48', side areas 33,33' of the end flange 32 and side plates 60 of the holding member 24.

10           The modified sample chamber 20 (and/or holding member 24) allows for the deposition of a cell substrate 56 at a specified reproducible location on a slide. The location is chosen such that the cell pellet 56 deposited by the modified sample chamber 20  
15 can function as a substrate in an overlay assay.

          The determination of the point on the slide on which to deposit the cell substrate is determined empirically as it will differ slightly for each individual rotating platform (rotating table, for  
20 example a Labline Orbit Shaker) due to mechanical differences as well as for the cell types, cell density, slide angle, rotational speed, temperature and other factors known to those skilled in the art. The determination can be made by depositing a test  
25 cell suspension on a slide that is on the rotating platform. The slide is then observed under the adherence assay test conditions and the area where the

0935016 072199  
66720 912550

cells aggregate is identified. The distances A and A' and the distance B from the center of the slide is determined for the area of cell aggregation as shown in Figure 17. A sample chamber 20 is then modified as described herein above to deposit the cytocentrifuge cell pellet substrate 56 at the calculated position as shown in Figure 18. Alternatively, a series of incrementally modified sample chambers 20 can be used to establish test slides and the modified sample chamber 20 that provides a cytocentrifuge cell pellet substrate 56 with the most appropriate positioning under the shear conditions established for the given adherence assay can be used. In Example 1 a protocol for an assay to test L-selectin ligand adhesion interactions is described, however other adherence assays requiring shear forces can also be undertaken with the present invention.

Selectins are involved in a variety of adhesive interactions, with biological roles ranging from acute inflammation (regulation of leukocyte trafficking) to immune responses (activation of T cells and regulation of lymphocyte migration to lymphoid tissues) to hematopoiesis. Studies by applicant on the interaction of L-selectin and ligands on hematopoietic cells which function as an adhesive receptor-ligand pair, i.e. *in vitro* binding studies of lymphocytes to KG1a, a primitive CD34-positive human

cell line derived from an acute myeloid leukemia (Civin et al., 1994; Koeffler et al., 1980) lead to the present invention. These studies surprisingly revealed highly specific adherence of lymphocytes to KG1a cells mediated by L-selectin on the lymphocyte, but unexpectedly not involving CD34 as the corresponding ligand as had been previously reported (Baumhueter et al, 1993). The results indicated the presence of a ligand, designated SILL (the ligand has also been referred to as hematopoietic cell L-selectin ligand, HLL), for L-selectin on the surface of this hematopoietic progenitor cell line and provide the first evidence of sulfation-independent L-selectin-mediated adhesion between lymphocytes and a non-endothelial cell type.

For these studies, the lymphocyte-HEV adherence assay was used which is an *in vitro* approximation of physiologic adhesion mediated by L-selectin was used and as discussed herein above is the standard assay for determining adhesion. This assay required the functional interaction of respective membrane structures localized within natural cell ligand bilayers. It has been a fundamental tool in studying the function of L-selectin in its native state on the surface of lymphocytes. This binding assay was novelly adapted as described herein (Figures 10-18) to examine lymphocyte-hematopoietic cell

adhesion, and the results provide the unexpected results of L-selectin-dependent adhesive interactions between lymphocytes and non-endothelial, hematopoietic cells.

5           Blood cell formation depends critically upon discrete cell-cell and cell-matrix adhesive receptor/ligand interactions which create specialized bone marrow microenvironments wherein cells proliferate and mature. L-selectin is  
10   characteristically expressed on a subpopulation of bone marrow cells which includes the earliest multilineage hematopoietic stem cells. Evidence that L-selectin may function in hematopoiesis has been  
15   obtained from both *in vitro* clonogenic assays (Koenig et al., 1994) and from clinical engraftment data following bone marrow transplantation (Dercksen et al., 1995).

          L-selectin ligands have been recognized heretofore only on endothelial cells and the function  
20   of all have been shown to be sulfation dependent (see United States Patent 5,489,578). The detection of a sulfation-independent L-selectin ligand on a non-endothelial cell (Example 6) expands the physiologic implications of L-selectin function beyond its well-  
25   characterized role in regulating leukocyte trafficking.

09363416-07216

The adaptation of the assay allowed for the first time the use of cell lines in a lymphocyte-HEV adherence assay. In this assay, slides were prepared utilizing a modified cytocentrifugation apparatus (Figures 10-18) of KG1a cell suspensions which were used in place of slides of frozen lymph node sections as taught by the prior art. The KG1a cells were placed on the slides by using cytocentrifugation as further described hereinbelow in Example 1.

Several independent lines of evidence indicate that lymphocyte binding to KG1a is mediated primarily, if not solely, by L-selectin. First, an anti-L-selectin mAb (LAM1-3) previously shown to block L-selectin-mediated adherence to LN HEV (Spertini et al., 1991), completely inhibited PBL from binding to KG1a or LN HEV, whereas anti-CD45 and isotype control antibodies did not block lymphocyte binding. Second, L-selectin-mediated binding is a calcium-dependent event, and lymphocytes were unable to bind to KG1a in the presence of the calcium chelator EDTA. Third, carbohydrates such as man-6-P, PPME, and fucoidin inhibited lymphocyte adherence to KG1a. These compounds are all known to bind to L-selectin and to inhibit lymphocyte binding to HEV in the *in vitro* assay (Stoolman and Rosen, 1983; Stoolman et al., 1984). Lastly, it is known that PMA treatment of lymphocytes causes shedding of membrane L-selectin via

02546-02199  
661220-97185260

a protein kinase C activation pathway, and corresponds to the loss of lymphocyte binding to LN HEV in the *in vitro* assay (Tedder et al., 1990). In these studies, PMA-treated PBL were no longer able to bind to KG1a.

5           These results revealed that KG1a possess a high affinity L-selectin ligand, however, a significant role for CD34 in this ligand activity was excluded by the following observations: (1) treatment of KG1a cells with a variety of anti-CD34 monoclonal  
10 antibodies, singly and in combination, had no effect on binding; (2) the CD34-positive human hematopoietic cell lines RPMI 8402 and MO7e did not support lymphocyte adherence despite higher levels of CD34  
15 expression than that of KG1a; and (3) KG1a cells separated into subpopulations expressing high and low levels of CD34 showed equivalent binding (Oxley and Sackstein, 1994b). Treatment of KG1a with the enzymes neuraminidase, chymotrypsin, and bromelain abrogated  
20 lymphocyte binding to the cells, indicating that the ligand is a glycoprotein and that expression of ligand sialic acid is critical for function (Oxley and Sackstein, 1994b).

          The nature of the ligand was investigated by determining the effects of various enzyme treatments  
25 of KG1a on the binding capacity. Previous studies have shown that ligand expression of sialic acid is essential for L-selectin-mediated binding of

0152.00344-0152.00344



lymphocytes to LN HEV (Rosen et al., 1985). In the present examples, neuraminidase-treated KG1a showed a complete loss of lymphocyte binding, indicating that sialic acid residues are also a necessary component on the KG1a L-selectin ligand; as such, lymphocyte adherence to KG1a involves carbohydrate motifs and is not based strictly on protein-protein interactions. This finding, combined with the results of protease experiments, indicates that the KG1a ligand is a glycoprotein.

To examine whether O-linked glycosylations on the ligand play a central role in the adhesive interaction, KG1a were digested with the enzyme O-sialoglycoprotein endopeptidase which specifically cleaves proteins at sites of O-linked sialoglycosylation (Abdullah et al., 1992) and which has been shown to differentially cleave epitopes of the CD34 molecule (Sutherland et al., 1992). The data revealed that treatment of KG1a in suspension with the enzyme actively destroyed CD34 epitopes, yet had no effect on lymphocyte adherence. These results suggest that ligand sialic acid residues critical to binding are present on N-linked rather than on O-linked glycosylations.

The functional dependence on intact sialic acid moieties is consistent with results from endothelial L-selectin ligands (True et al., 1990) and

0935846-072199

667220" 918560  
suggested that epitopes critical for binding might be structurally conserved among these ligands. However, treatment of KG1a with O-sialoglycoprotease resulted in characteristic loss of enzyme-sensitive epitopes  
5 from CD34 (Sutherland et al., 1992) but did not affect adherence (Oxley and Sackstein, 1994b). Since the ligand activity of endothelial CD34 is eliminated by this treatment (Puri et al., 1995), this result provided further evidence that the KG1a ligand  
10 activity was not attributable to CD34, and, more generally, that the binding determinant of the ligand differs structurally from proteins recognized by MECA-79.

CD34 has been reported to be a ligand for L-  
15 selectin based on the finding that a murine L-selectin-IgG chimera molecule precipitated CD34 from a murine lymph node lysate (Baumhueter et al., 1993). The results as set forth in the examples indicate that CD34 as expressed on KG1a is not a functional ligand  
20 for lymphocyte L-selectin, as no difference in lymphocyte binding to sorted CD34- and CD34+ KG1a cells was observed (see Figures 3 and 4). Titration studies using varying proportions of KG1a and HL60 have demonstrated that the amount of lymphocyte  
25 adherence is directly proportional to the percentage of input KG1a cells, indicating that differences in lymphocyte binding to the positive and negative sorted

fractions would have been evident if CD34 were the ligand. It is unlikely that a particular binding epitope of CD34 as selected, as this experiment was done using two different anti-CD34 mAbs to sort the KG1a. Two forms of CD34 on KG1a have been reported ("truncated" and "full length") (Krause et al., 1993); however, these differences do not account for the data herein as sorting was also performed using QBEND10, which recognizes both forms.

10 In addition to sorting experiments, evidence that CD34 is not the L-selectin ligand on KG1a is derived from mAb blocking studies and adherence assays using other CD34 positive cells. None of the anti-CD34 mAbs tested, or any combination thereof, was able to block lymphocyte binding to KG1a. Furthermore, lymphocytes did not adhere to another primitive CD34+ cell line, RPMI 8402, and transfection of CD34 into COS-7 cells did not confer lymphocyte binding capacity. While potential glycosylation differences of the CD34 molecule expressed by these cell types could affect their ability to support lymphocyte adherence, this explanation is unlikely in light of equivalent adherence observed among the sorted CD34+ and CD34- KG1a cells. Taken together, the data presented here indicate that the CD34 glycoform present on hematopoietic cells is not a ligand for L-selectin. Moreover, flow cytometric analysis of the

various cell lines utilized in the binding assay provides evidence that membrane structures such as LFA-1, VLA-4, CD44, Sialyl Le<sup>x</sup> and CD43 do not play a primary role in lymphocyte adherence to KG1a since  
5 each of these molecules were also present on at least one other cell line tested that did not demonstrate lymphocyte binding.

SILL is not recognized by MECA-79 monoclonal antibody (Example 5) which identifies L-selectin  
10 ligands on lymph node HEV. Immunofluorescence analysis of KG1a using MECA-79 shows no evidence of the protein identified by MECA-79. SILL is shown to be unique from L-selectin ligands thus far identified.

The relationship of the KG1a ligand to  
15 proteins identified by MECA-79 were evaluated by performing adherence studies in the presence of MECA-79, flow cytometric analysis of MECA-79 antigens on KG1a, and immunoprecipitation studies of metabolically radiolabeled KG1a and lymph node organ cultures.  
20 Incubation of KG1a with MECA-79 did not interfere with lymphocyte adherence in the binding assay and flow cytometric studies showed no evidence of MECA-79 antigen on KG1a (Figure 6A-B). Moreover, no MECA-79-immunoprecipitable proteins were detected from lysates  
25 of metabolically radiolabeled KG1a cells using either <sup>35</sup>SO<sub>4</sub> (Figure 6C) or <sup>35</sup>S-methionine/cysteine, though characteristic bands were obtained from respectively

093546-0419  
66720-978560

radiolabeled murine lymph nodes (Figure 6C). The absence of MECA-79-precipitable protein within KG1a lysates indicates that there are no intracellular accumulations of MECA-79 antigen, and, furthermore, that the MECA-79 antigen is not synthesized by the KG1a cells.

Direct cell-cell interactions were utilized to detect the presence of an L-selectin ligand on a hematopoietic cell. Other studies directed at identifying L-selectin ligands have relied on molecular approaches utilizing a murine L-selectin-IgG chimera molecule, synthesized in a human embryonal kidney cell line, as a probe (Watson et al., 1990). Of note, studies utilizing this chimera have failed to demonstrate binding of the molecule to KG1a cells (Majdic et al., 1994). In general, tissue- and species-specific patterns of glycosylations are well described, (Yamashita et al., 1983; Cullen et al., 1981; Yamashita et al., 1985) and such differences can affect the biological activity of proteins expressed in different cells (Cowing 1983; Huff et al., 1983). As it is known that glycosylation of L-selectin varies among different cells expressing the protein (Lewinsohn et al., 1987, Ord et al., 1990; Griffin et al., 1990), such differences may account for the observation here that native L-selectin, expressed on lymphocyte membranes, selectively binds to a

093516-072199

corresponding ligand on KG1a cells while the chimera  
apparently does not. Similarly, differences in  
glycosylation of CD34 among endothelial cells and  
hematopoietic cells may account for the differential  
5 capacity of this protein to participate in L-selectin  
interactions among these cell types.

Recognition of endothelial ligands by MECA-  
79 is sulfation-dependent, however, absence of MECA-79  
reactivity does not in itself indicate that sulfation  
10 does not participate in the binding determinant of the  
KG1a ligand of the present invention. Applicant  
directly examined this by measuring ligand activity  
following incubation with chlorate (Example 6), a  
highly efficient metabolic inhibitor of both  
15 saccharide and peptide sulfation (Baeuerle and  
Huttner, 1986). The results of chlorate incubation on  
L-selectin ligand activity of KG1a was evaluated and  
the results are shown herein below (Table 3 and Figure  
9). Incubation of KG1a in chlorate-containing media  
20 was performed following complete desialylation of the  
cells by neuraminidase treatment. Initial studies of  
the kinetics of recovery of binding activity following  
either neuraminidase treatment or protease digestion  
of membrane protein showed absence of ligand activity  
25 for twelve hours; thereafter, ligand activity  
increased in a linear fashion with return to baseline  
levels within twenty-four hours. Return of binding

09350116.072199  
66720.912560

activity following desialylation or protease treatment was blocked by metabolic inhibition of N-linked glycosylation by tunicamycin treatment, suggesting that re-expression of activity results predominantly from synthesis/processing of nascent ligand and not from transport to membrane of stored mature protein (Table 3).

Accordingly, KG1a were incubated in chlorate for twenty-four hours to inhibit sulfation throughout the duration of ligand reexpression.  $^{35}\text{SO}_4$ -pulse radiolabeling studies at eight hour time intervals within the twenty-four hours of chlorate incubation indicated that sulfation was inhibited throughout the entire incubation period, as demonstrated by trichloroacetic acid-precipitable radiolabeled protein counts (chlorate-incubated counts consistently <10% that of non-chlorate treated controls) and SDS-PAGE analysis of radiolabeled protein of cell lysates. As shown in Table 3, the binding activity of the KG1a ligand was unaffected by chlorate incubation, though inhibition of sulfation was confirmed, even for the terminal six hours of the incubation period, by a marked reduction of  $^{35}\text{SO}_4$  incorporation into CD43, a sulfated sialomucin (Wilson and Rider, 1992) expressed on KG1a (Oxley and Sackstein, 1994b) (see Figure 9). The data shows that sulfation, regardless of whether it may be present on protein or sugar, is not critical

0935016 072199

to the functional interaction with L-selectin. These data, together with the lack of MECA-79 reactivity, indicate that sulfation does not contribute significantly to the function of the KG1a ligand.

5           The data presented herein expands the understanding of the structure of selectin ligands and provides the first evidence of a functional, membrane glycoprotein L-selectin ligand whose binding activity is not sulfate-dependent. Multiple studies have  
10 emphasized the importance of sulfation in the binding determinant of ligands for both L-selectin (Hemmerich et al., 1994; Imai et al., 1993; Suzuki et al., 1993; Green et al., 1995; Stoolman and Rosen, 1983; Green et al., 1992; Hemmerich and Rosen, 1994b) and P-selectin  
15 (Aruffo et al., 1991; Sako et al., 1995; Pouyani et al., 1995), and all previously identified naturally-occurring membrane ligands for these selectins bear sulfate-dependent activity. The contribution of sulfate modifications may relate to the localization  
20 of negative ions, on either sugar (Hemmerich and Rosen, 1994b) or protein (Sako et al., 1995; Pouyani, 1995; Niehrs et al., 1992) determinants. Such a role for charge is supported by the finding that unsulfated anionic polysaccharides such as polymers of phosphated  
25 mannose can bind to L-selectin (Yednock et al., 1987). Within the KG1a ligand, it is possible that glycosidic and/or amino acid modifications such as

0152.00344



hosphorylation, or the molecular composition of the discrete sugars or amino acids comprising the binding domain, create a relevant anionic milieu. Although the precise structural features which direct binding activity for this and other L-selectin ligands remain to be determined, that data indicate that cell membrane binding determinants conferring high affinity recognition of L-selectin are not strictly conserved: while sulfated glycoprotein ligands may be characteristic of endothelial cells from lymphoid tissues, non-sulfated ligand forms may direct L-selectin-specific cell-cell recognition and adhesion events in other cell types, be it non-endothelial cells or non-lymphoid endothelium.

The above discussion provides a factual basis for the characterization and use of SILL and the method and apparatus for performing adherence assays.

The methods used with and the utility of the present invention can be shown by the following examples.

## EXAMPLES

### GENERAL METHODS:

*Cell Lines.* Cell lines used in these studies were obtained from the following sources: KG1a and Nalm 16, gift of Dr. William E. Janssen; HL60, K562, and Raji, gift of Dr. Lynn Moscinski; COS-7, gift of Dr. Kenneth Zukerman (all from H. Lee Moffitt

Cancer Center, Tampa, FL); RPMI 8402, gift of Dr. Daniel G. Tenen (Harvard Medical School, Boston, MA).

All cells were cultured in RPMI 1640 (Gibco-BRL, Gaithersburg, MD) supplemented with 10% heat-inactivated fetal bovine serum (FBS) in a humidified chamber at 37°C with 5% CO<sub>2</sub> in air.

*MECA-79 antibody.* MECA-79 antibody was a gift from Dr. Phillip Streeter, Searle Research Laboratories/Monsanto Co., St. Louis, MO). A MECA-79 hybridoma is available from American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, MD 20852 as accession number HB-9479. The production of the MECA-79 hybridoma is described in United States Patent 5,403,919.

*Preparation of Lymphocytes.* Human peripheral blood lymphocytes (PBL) were isolated by Ficoll density gradient from blood drawn in sodium citrate. To obtain rat thoracic duct lymphocytes (TDL), thoracic ducts of rats were cannulated as described by Bollman et al. (1948). Lymph was collected in phosphate buffered saline (PBS) with 0.1% penicillin/streptomycin and 5 U/ml heparin. PBL or TDL were washed three times in RPMI 1640 medium without bicarbonate (Gibco-BRL), pH 7.4, and suspended at  $1 \times 10^7$  cells/ml in above medium with 5% FBS and kept on ice until use in the adherence assay.

5 *Lymphocyte Adherence Assay.* The procedure  
for the *in vitro* binding of human or rat lymphocytes  
to KG1a was adapted from the rat lymphocyte-lymph node  
binding assay which has been described by Stamper and  
10 Woodruff (1976) and Sackstein et al. (1988) and  
described in Example 1. Cytocentrifuge preparations  
of KG1a or other cell lines were made on a Cytospin 3  
Cytocentrifuge (Shandon Lipshaw, Pittsburgh, PA).  
Frozen rat LN sections 8 $\mu$ m thick were mounted on  
15 slides, and lymphocyte binding to LN HEV served as a  
positive control in all experiments. Slides were air  
dried, fixed in 3% glutaraldehyde (Electron Microscopy  
Sciences, Fort Washington, PA) in PBS, rinsed with  
PBS, incubated in 0.2M L-lysine (Sigma Chemical  
20 Company, St. Louis, MO) to block unreacted  
glutaraldehyde, then rinsed and held in RPMI 1640 with  
1% FBS at 4°C until use in experiments.

Lymphocyte suspensions (200  $\mu$ l) were  
overlaid onto cytocentrifuge or LN sections in  
20 duplicate and placed on a rotating platform (80 rpm)  
at 4°C for 30 minutes. Slides were then rinsed in cold  
PBS to remove non-adherent lymphocytes, fixed in 3%  
glutaraldehyde, and stained with methyl green-thionin.  
Slides were examined under the light microscope for  
25 adherence of lymphocytes to KG1a or LN HEV.

09358115-072199

Number of lymphocytes adherent to confluent area of KG1a were counted by light microscopy using an ocular grid under 250× magnification. Quantitation was performed by examining two fields per slide, minimum of two slides per experiment, three separate experiments. Results are presented as % binding compared to corresponding untreated control sections.

*Treatment of Lymphocytes with Potential Inhibitors.* Lymphocytes in RPMI 1640 medium with 5% FBS were pre-incubated (30 minutes on ice) and the assay performed in the presence of the following inhibitors: 1 mM EDTA (no pre-incubation period); 10 mM D-mannose-6-phosphate (Sigma); 10 µg/ml PPME (kindly provided by Dr. M.E. Slodki, USDA, Peoria, IL); and 5 µg/ml fucoidin (Sigma).

*Antibody Blocking Experiments.* Lymphocytes ( $1 \times 10^7$  cells/ml) were pre-incubated on ice for twenty minutes with mAbs at 1.0 µg/ml and used in the binding assay without further washing. The following mAbs were used: LAM1-3 (anti-L-selectin) (kind gift of Dr. Thomas Tedder, Duke University, Durham, N.C., and also obtained from Coulter Corp., Hialeah, FL); anti-CD45 (leukocyte Common Antigen) (Becton Dickinson, San Jose, CA); and IgG<sub>1</sub> (isotype control) (Coulter). In some experiments, prepared KG1a slides were incubated with 0.2 µg of anti-CD34 antibodies {HCPAa-1 (clone

My10) and HPCA-2 (clone 8G12) (Becton Dickinson), QBEND10 (AMAC) and 12.8 (kindly provided by Dr. Pat Roth, Coulter Corp.)} in RPMI 1640 with 5% FBS for 30 minutes prior to the binding assay.

5            *PMA Treatment of Lymphocytes.* Lymphocytes were suspended at  $1 \times 10^7$  cells/ml in cell culture medium and incubated 1 hour at 37°C with or without 10 ng/ml PMA (Gibco-BRL). Cells were then washed twice in PBS and used in either the lymphocyte binding assay  
10 or analyzed for surface antigens by flow cytometry (see below).

*Enzyme Treatment of KG1a or LN.*

Cytocentrifuge preparations of KG1a or LN frozen sections were glutaraldehyde-fixed, then treated with  
15 various enzymes prior to the binding assay. For treatment with neuraminidase (sialidase), slides were rinsed twice with enzyme buffer (50 mM NaAc, 154 mM NaCl, 9 mM  $\text{CaCl}_2$ , pH 5.5), then incubated 30 minutes at 37°C with 50  $\mu\text{l}$  of buffer (control) or undiluted  
20 neuraminidase (1.2 U/ml, Boehringer Mannheim, Indianapolis, IN). In protease studies, slides were incubated with RPMI 1640 alone or RPMI 1640 containing enzymes: 100 U/ml chymotrypsin (Sigma) (115 minutes at 37°C), or 0.1% bromelain (Sigma) (30 minutes at  
25 37°C); to assess specificity, the protease inhibitors PMSF (1.0 mg/ml, Sigma) and chymostatin (900  $\mu\text{g/ml}$ ,

00344-0219

Boehringer Mannheim) were coincubated with  
chymotrypsin (100 U/ml) for 15 minutes at 37°C.

Following enzyme treatments, slides were washed three  
times with RPMI 1640 and placed in RPMI 1640 with 1%  
5 FBS until use in the binding assay.

KG1a cells in suspension ( $4 \times 10^7$  cells/ml)  
were incubated with O-sialoglycoprotein endopeptidase  
(Accurate Chemical and Scientific Corp., Westbury, NY)  
(0.24 mg/ml, 37° C, 30 minutes), washed three times  
10 with 2% FBS in PBS, and cytocentrifuge preparations  
were made for use in the binding assay. To verify the  
activity of the enzyme, cells were tested for the  
cleavage of CD34 by flow cytometry using QBEND10 mAb.

*Antigen Expression by Flow Cytometry.* Flow  
15 cytometric analysis was performed using the following  
commercially-available mAbs together with isotype-  
matched controls: TQ1 (anti-L-selectin), LAM1-3  
(Anti-L-selectin), 4B4 (anti-VLA-4) (all from Coulter  
Corp.); QBEND10 (anti-CD34) (AMAC, Westbrook, ME);  
20 anti-CD44, LFA-1- $\beta$  (anti-CD18), LFA-1- $\alpha$  (anti-CD11a),  
HPCAS-2 (anti-CD34), anti-CD45, Leukosialin (anti-  
CD43), anti-Sialyl-Le<sup>x</sup> (all from Becton Dickinson).  
Cells ( $1 \times 10^6$ ) in 100  $\mu$ l of PBS with 2% FBS were  
incubated on ice for 25 minutes with antibody as per  
25 manufacturer's recommendations, washed  $\times 3$  and analyzed  
on a FACStar<sup>PLUS</sup> (Becton Dickinson).

035515-0219  
66720-918560

Fluorescence Activated Cell Sorting of KG1a cells. KG1a cells were stained with anti-CD34 mAbs (QBEND10-FITC in two experiments, HPCA-2-PE in one experiment) and positive and negative expressing cells were sorted on a FACstar<sup>PLUS</sup> flow cytometer equipped with an argon laser tuned at 488 nm (Becton Dickinson). Sorted cell populations were restained with anti-CD34 antibody directed at epitopes not used for sorting and were analyzed to determine the efficiency of the sort. Cytocentrifuge preparations were made of the positive and negative sorted fractions and were used in the lymphocyte binding assay.

*Transfection of COS-7 with CD34 cDNA.* COS-7 cells were transiently transfected with human full-length CD34 cDNA in pCDM8 plasmid (a gift from Dr. Daniel Tenen, Boston, MA) using a DEAE Dextran transfection method (Selden, 1992). Briefly, COS-7 cells were incubated for 4 hours at 37°C with 10 ml of transfection solution containing 20-40 µg of plasmid DNA, 10% Nu Serum (Collaborative Biomedical Products, Bedford, MA), 400 µg/ml DEAE Dextran (Sigma), and 100 µ chloroquine (Sigma) in Dulbecco's Modified Eagles Medium (Gibcon-BRL). Cells were then rinsed and treated with 10% DMSO (Sigma) in PBS for two minutes at room temperature, rinsed in PBS, and incubated in

0152.00344

003316-07219  
66720-912560

tissue culture media for three days. In one set of experiments, trypsinization was avoided by growing transfected cells directly on glass slides for subsequent use in the binding assay or for analysis of CD34 expression by fluorescence microscopy. In other experiments, COS-7 cells grown on 10 cm plates were removed with trypsin/EDTA (0.25%/1 mM, Gibco-BRL), then analyzed for CD34 expression by flow cytometry. These trypsinized cells were then placed on slides by cytocentrifuge for use in the lymphocyte binding assay.

*Flow Cytometric Analysis of KG1a and Immunoprecipitation of <sup>35</sup>S-metabolically radiolabeled KG1a.* Indirect immunofluorescence staining was performed by incubating  $1 \times 10^6$  KG1a with 1  $\mu$ g MECA-79 or IgM control antibody (both gifts from Dr. Phillip Streeter, Searle Research Laboratories/Monsanto Co., St. Louis, MO) in 2% FBS in PBS for 30 minutes on ice.

After washing, the cells were treated with 5  $\mu$ g FITC-conjugated secondary antibody for 30 minutes on ice. Flow cytometric analysis was performed on a Facscan flow cytometer (Becton-Dickinson). KG1a were metabolically radiolabeled by incubating  $100 \times 10^6$  cells ( $20 \times 10^6$  cells/ml) for five hours in CRCM30 sulfate-free medium (Sigma Chemical Co., St. Louis, MO) supplemented with 150  $\mu$ Ci/ml <sup>35</sup>S-Na<sub>2</sub>SO<sub>4</sub> (Dupont/New



England Nuclear, Boston, MA). Mesenteric lymph nodes from three mice (~60 mg of tissue) were minced and radiolabeled in parallel with the KG1a for five hours in CRCM30 sulfate-free medium supplemented with 1mCi/mi  $^{35}\text{S-Na}_2\text{SO}_4$ . KG1a and murine lymph nodes were lysed in 2% Triton X-100 in Tris-buffered saline (TBS) containing 1 mM PMSF and 1  $\mu\text{g/ml}$  each of aprotinin, leupeptin and pepstatin A overnight at 4°C. The lysates were clarified by centrifugation at 10,000g for fifteen minutes and the samples precleared with protein G-agarose for four hours. Immunoprecipitation was performed using equivalent amounts of TCA-precipitable counts from lymph node and KG1a lysates, using 2  $\mu\text{g}$  IgM control antibody or MECA79 followed by 6  $\mu\text{g}$  goat anti-rat IgM secondary antibody (Jackson ImmunoResearch Labs Inc.), or 1  $\mu\text{g}$  anti-CD43 or IgG<sub>1</sub>, followed by protein G agarose overnight incubation at 4°C. Samples were electrophoresed (7.5% SDS-PAGE) under reduced conditions followed by autoradiography as described previously (Sackstein et al., 1995).

*Chlorate Inhibition of Sulfation.* The KG1a adherence assay was performed and quantitated as described previously (Oxley and Sackstein, 1994b). The anti-CD43 antibody L60 (Becton-Dickinson) recognizes a sialylated epitope, thus allowing an independent assessment of the efficiency of

093516-0319

neuraminidase treatment of KG1a. KG1a were treated with 0.1 U/ml neuraminidase (*V. cholerae*, Boehringer Mannheim, Indianapolis, IN) for one hour in RPMI 1640 without bicarbonate (Gibco/BRL, Gaithersburg, MD) for  
5 sixty minutes at 37°C. Control cells were incubated in RPMI 1640 without bicarbonate with an equivalent volume of neuraminidase storage buffer. The cells were washed in RPMI 1640 twice, tested by the adherence assay to verify complete loss ligand  
10 activity, and by flow cytometry to verify loss of the L60 epitope. The neuraminidase-treated KG1a were recultured in RPMI 1640 containing 10% FBS ( $2 \times 10^6$  cells/ml) in the presence or absence of 10 mM sodium chlorate (Sigma Chemical Co.) for twenty-four hours.  
15 To verify chlorate inhibition of sulfation, CD43 was immunoprecipitated from equivalent numbers of untreated and chlorate-treated KG1a that were  $^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$  radiolabeled for the last six hours of the twenty-four hour chlorate treatment.

20

#### EXAMPLE 1

The procedure for the *in vitro* binding of human or rat lymphocytes to KG1a cells was adapted  
25 from the rat lymphocyte - lymph node (frozen section) binding assay which has been described by Stamper and Woodruff (1976) and Sackstein et al. (1988) as

09353116-072199

described herein above. Cytocentrifuge preparations of KG1a or other cell lines were made on a Cytospin 3 Cytocentrifuge (Shandon Lipshaw, Pittsburgh, PA) following the manufacturer's instruction (Cytospin®3  
5 Cell Preparation System Operator Guide) with modifications of the sample chamber to provide appropriate placement of the cytocentrifuge cell pellet onto slides for use in an adherence assay as described herein above.

10                    Cell Separation: Cell separation procedures for blood cells as are generally known to those skilled in the art are used to separate lymphocytes and granulocytes from peripheral blood samples, from bone marrow aspirates, and from vertebral bodies  
15 donated through organ harvests. More specifically:

Peripheral Blood

*Sub B1* (1) Sodium citrated blue top tubes are used, spun approximately 30 seconds at 100g to pull all cells away from the top of the tubes. The blood  
20 is removed, 25 ml or less, and placed into a 50 ml Falcon® tube. Using sterile 1M PBS, add equal volume to the volume of blood for a 1:1 final dilution.

(2) Place 10 ml of FICOLL lymphocyte separation medium into a fresh 50 ml tube. Layer the  
25 blood from step 1 onto the top of the FICOLL medium slowly and carefully to insure the interface is tight. Place no more than 35 ml of diluted blood on top of

0152.00344

the FICOLL layer so that the total volume inside the tube does not exceed 45 ml. Spin at 400g for 30 minutes.

(3) Remove the upper clear layer  
5 containing the plasma and platelets; remove the next layer (cloudy, lymphocyte layer, PBL: peripheral blood lymphocytes), and place in a 50 ml tube. Suspend the PBL to 45 ml in 1x PBS. Spin them again for 10 minutes at slightly less than 200g (any more force  
10 than this will cause platelet contamination). Count cells, wash and spin at 1000 rpm twice and resuspend at 3 to 4 million cells per ml which is the volume necessary to achieve correct cell density in the cytocentrifuge procedure.

(4) Remove the clear FICOLL layer from  
15 the tubes from step (3) and discard leaving only the granulocytes and monocytes in the red cell pellet. Dilute the pellet 1:1 by volume with 1x Hank's Balanced Salt Solution (HBSS). Dilute this volume 1:1  
20 with 3% Dextran (made by dissolving 1.5g of Dextran powder into 50 ml of sterile 1x PBS) and shake gently to mix. Allow to stand for 25 minutes so that the red blood cells pellet out under gravity, leaving the granulocytes and monocytes in the supernatant. After  
25 25 minutes, remove the supernatant and dilute it to 45 ml in 1x PBS. Discard the red cell pellet. Spin the

093516-0249  
66720-918560

granulocytes and monocytes again for 10 minutes at 200g. To remove remaining red cells treat pellet with a salt gradient. Count the cells and wash one more time at 200g for 10 minutes before resuspending to 3 to 4 million cells per ml. Both groups of cells (PBL and granulocytes) can be held on ice until ready for cytocentrifuge.

Sub  
Ba  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
10230  
10235  
10240  
10245  
10250  
10255  
10260  
10265  
10270

remember the danger associated with working with patient blood. Due to the nature of the procedure used to retrieve the desired cell populations, special precautions must be taken to prevent contact with the fluids and tissues of the sample by the researcher. A full frontal covering must be worn along with gloves and a face guard.

(1) Cut the vertebral body in half and remove the marrow and placed into a solution of heparin. It is better to remove many very small pieces of marrow instead of just a few large chunks. This allows for more surface area of marrow to be exposed, and therefore, more cells can be collected from the bone marrow matrix.

(2) Once all of the marrow has been removed, shake the tube containing the heparinized marrow vigorously, in order to free any cells attached to the matrix. Let the tube stand for a few seconds to allow all of the dense marrow material to sink to the bottom of the tube. Remove the supernatant solution and place it into a clean tube. Resuspend the chunks of marrow in heparin again and shake vigorously to free more cells. Again, remove the supernatant. Repeat several times. Once the marrow is cleansed of most of the adherent cells, spin the supernatant materials in the centrifuge for 10 minutes at 200g. Resuspend the pellet in RPMI medium and pass

the suspension through a sterile nylon mesh filter into a clean 50 ml tube. This will remove any remaining marrow pieces. The remaining cells should be counted and washed two more times before being  
5 suspended to proper concentration (3 to 4 million per ml) in RPMI medium for cytocentrifuge.

After each of the cell populations has been separated, washed, and suspended to proper concentration, the following procedure is used to  
10 place the cells onto slides in order to test the binding capacity of each population. The cytocentrifuge will pull cells out of small volumes of solution and place them one cell layer thick onto slides. If the concentration of cells in solution is  
15 too high, cells will be stacked on top of one another and the experiment will not produce useful results. If the concentration is too low, the actual numerical data collected from the experiment will not be statistically valuable. Cell suspensions should have  
20 a concentration of only 3 to 4 million cells.

#### Cytocentrifuge Procedure

(1) As described herein above, place a cell suspension on a slide and place on the rotational  
25 table under the experimental conditions of the adherence assay. Observe where cells aggregate and prepare a model slide (Figure 17) showing this

0152.00344-0249

location by which to mark all other slides.

(2) After marking the proper number of slides (i.e. typically one slide/ $2.5 \times 10^5$  cells), take the same number of clamps, funnels, and filter cards (paper guides) (Figure 10) as slides needed. Assemble the holder assembly and load in cytocentrifuge as per the instructions in the cytocentrifuge manual. Once the holder assemblies are loaded, place 100  $\mu$ l of the proper cell suspension in each funnel. Set speed at 570 and time 5 minutes for this particular assay. As per instruction in manual remove slides from holding member after spin. Do not smear or wipe the delicately placed cells from the slide. Allow slides to dry for one minute.

(3) When dry, place the slides into a 3% glutaraldehyde solution (3.6 ml of 25% glutaraldehyde to 26.4 ml of 1x PBS per 12 slides) for ten minutes. After the 10 minutes of fixing, rinse the slides twice, gently, in 1x PBS. Place the slides into lysine solution (0.2M) for 10 minutes. This solution will prevent nonspecific binding in the adherence assay. After lysine treatment, rinse slides twice in RPMI and then place them into 1% FBS in RPMI at 4°C until ready to perform the adherence assay.



## EXAMPLE 2

*Lymphocytes Bind to KG1a.* Lymphocytes (both PBL and TDL) adhered specifically and reproducibly to KG1a, but not to RPMI 8402, HL60, Nalm 16, K562, or Raji cell lines in the *in vitro* binding assay (Table 1). All experiments were performed in parallel with LN frozen sections as positive controls. Lymphocyte binding to KG1a was observed under conditions identical to those whereby L-selectin mediates binding of lymphocytes to LN HEV.

*Lymphocyte Binding to KG1a is Mediated by L-selectin.* To directly examine whether lymphocyte attachment was mediated by L-selectin, PBL were pre-incubated with the anti-L-selectin mAb LAM1-3, anti-CD45, or IgG<sub>1</sub> isotype control antibodies. The LAM1-3 antibody completely inhibited lymphocyte binding to KG1a and LN control, while CD45 and isotype control mAbs did not affect binding (Fig. 1A & 1B). In order to quantify the relative amounts of antibody attachment to lymphocytes, antibody-treated lymphocytes were incubated with goat-anti-mouse FITC-conjugated secondary antibody and analyzed by flow cytometry. Although the amount of anti-CD45 antibody on lymphocytes was significantly greater than that of LAM1-3 as indicated by mean channel fluorescence (Fig. 2A-C), LAM1-3 alone blocked lymphocyte adherence to

0152.00344-0219

KG1a and LN HEV, indicating that this effect was specific and not secondary to charge or stearic alterations of the lymphocyte membrane.

*The Effect of Enzyme Treatment of KG1a on*  
5 *Lymphocyte Binding.* Pretreatment of both KG1a and LN control sections with neuraminidase (60 mU), chymotrypsin (100 U/ml) or bromelain (0.1%) prior to the binding assay abrogated binding of lymphocytes, while treatment with buffer or medium alone did not  
10 alter binding capacity. In addition, the effects of chymotrypsin were confirmed by coincubation with the protease inhibitors chymostatin and PMSF, which prevented chymotrypsin effects on lymphocyte binding.

However, pretreatment of KG1a with O-  
15 sialoglycoprotein endopeptidase had no effect on lymphocyte binding despite complete enzymatic removal of the CD34 epitope recognized by QBEND10 mAb (see Table 2).

*Lymphocyte Binding to KG1a is Calcium*  
20 *Dependent.* Lymphocyte binding to KG1a and to LN control sections was completely inhibited by the presence of EDTA, indicating a calcium requirement for lymphocyte-KG1a binding.

*Mannose-6-Phosphate, PPME, and Fucoidin*  
25 *Inhibit Lymphocyte Binding to KG1a.* The specificity of lymphocyte-KG1a binding was investigated by treating PBL or TDL with carbohydrate inhibitors of L-

035815-072499

selectin-HEV interactions prior to the adherence assay. Man-6-P (10 mM), PPME (10 µg/ml), and fucoidin (5 µg/ml) all inhibited lymphocyte binding to both KG1a and LN control sections (see Table 2).

5           *PMA Treatment of Lymphocytes Results in the Loss of Binding to KG1a.* PBL were incubated for 1 hour at 37°C with 10 ng/ml PMA, then used in the lymphocyte binding assay. PMA-treated PBL were unable to bind to either KG1a or LN HEV, while control PBL  
10 demonstrated high amounts of binding (see Table 2).

          Loss of surface L-selectin was assessed by flow cytometric analysis of TQ1 levels in control and PMA-treated PBL. PMA-treated lymphocytes showed a dramatic decrease in TQ1 mean channel fluorescence (to  
15 levels less than 10% of that of untreated cells) in three separate experiments. PMA-treated PBL were also analyzed for expression of CD44, LFA-1 (both α and β chains), and VLA-4, and expression of these adhesion molecules following PMA exposure was identical to  
20 expression on control PBL.

### EXAMPLE 3

#### *Pretreatment of KG1a with Anti-CD34*

25   *Antibodies Did Not Inhibit Adherence of Lymphocytes.*  
Cytocentrifuge preparations of KG1a were preincubated

09358115-072199  
16T220-9T8560

with anti-CD34 antibodies and the binding assay was performed in the presence of the antibodies (Table 2).

Monoclonal antibodies to four different CD34 epitopes were used alone or in combination, including the

5 clones My10, QBEND10, 8g12, and 12.8, in amounts ranging from 0.2 to 17 µg/slide. Anti-CD45

(irrelevant control) and IgG<sub>1</sub> (isotype control) antibodies were also tested. None of the anti-CD34 antibodies inhibited lymphocyte binding to KG1a, despite immunohistochemical evidence of extensive antibody binding to the glutaraldehyde-fixed KG1a sections.

*Other Surface Antigens on KG1a do not Appear to Mediate Binding.* The surface expression of several antigens on KG1a, RPMI 8402, HL60, Nalm 16, K562, and Raji was analyzed by flow cytometry (Table 1). LFA-1, FLA-4, CD44, Sialyl Le<sup>x</sup>, and CD43 were all expressed by KG1a and at least one other cell line that did not support lymphocyte adherence. Of note, although RPMI 8402 cells express CD34 at levels comparable to KG1a, there was no adherence of lymphocytes to these cells in the binding assay.

*CD34 Positive and Negative KG1a Cells Supported Equivalent Amounts of Lymphocyte Binding.*

25 CD34+ and CD34- KG1a cells were separated by fluorescence activated cell sorting and cytocentrifuge preparations of each population were made. The in

035816-07199

vitro adherence of lymphocytes was identical in the CD34+ and CD34- populations despite an enrichment of >90% and <10% CD34+ cells in the respective populations (Figs. 3A-D and 4; Table 2).

5

**EXAMPLE 4**

*CD34-Transfected COS-7 Cells Did Not Support Lymphocyte Adherence.* COS-7 cells were transfected with CD34 and tested in the *in vitro* binding assay, and both trypsinized and intact transfected COS-7 cells failed to support lymphocyte adherence. By flow cytometric analysis, transfected cells were approximately 60% positive for CD34 expression, and the mean channel fluorescence was greater than that of KG1a control cells (Fig. 5A-F). Intact, untrypsinized COS-7 cells transfected with CD34 also strongly expressed CD34 ( $\approx$ 90% positive as estimated by fluorescence microscopy).

20

**EXAMPLE 5**

*Flow cytometric analysis of KG1a and immunoprecipitation of <sup>35</sup>S-metabolically radiolabeled KG1a demonstrate an absence of MECA79 antigen.*

25

In functional studies, incubation of KG1a cytocentrifuge layers with MECA 79 or isotype-control

09358115-072199

035316-072199  
66T27"9T35E60

(IgM) rat monoclonal did not inhibit lymphocyte attachment in the adherence assay (e.g., for studies using MECA 79 at a concentration of 50  $\mu$ g/ml, binding was 98.2%  $\pm$  3.38% (mean  $\pm$  SEM) that of control assays without antibody addition). However, as is characteristic of MECA 79, lymphocyte attachment to murine lymph node HEV was blocked using MECA 79 at concentrations as low as 2  $\mu$ g/ml. To investigate surface expression of MECA 79 antigen on KG1a cells, flow cytometric analysis of MECA-79 antigen was performed on KG1a cells and compared with isotype matched control antibody (Figs. 6A and 6B). Although lymph node HEV were clearly and characteristically identified by staining with MECA 79 as evident by fluorescence microscopy. There was no MECA 79 staining evident by flow cytometric analysis of KG1a cells by fluorescence microscopy.

To further evaluate for expression of MECA 79 antigens by KG1a, the cells were incubated with either  $^{35}$ S- $\text{Na}_2\text{SO}_4$  or  $^{35}$ S-methionine/cysteine and resultant radiolabeled protein was immunoprecipitated using MECA 79 antibody (Figure 6C). As a positive control, murine lymph nodes were radiolabeled, lysed, and immunoprecipitated in parallel with KG1a. To adjust for potential differences in efficiency of radioactive label incorporation between KG1a and murine lymph nodes, the relative amounts of lysate

immunoprecipitated from each source were equalized for  
TCA-precipitable counts. A representative  
immunoprecipitation of MECA 79 from  $^{35}\text{S}$ - $\text{Na}_2\text{SO}_4$ -labeled  
KGLa and lymph node is shown in Figure 6C. The  
5 characteristic 50,000 and 90,000 mw bands are  
immunoprecipitated by MECA 79 in lymph node lysate but  
no MECA 79 bands are detected in the KGLa lysate;  
similar results were obtained with MECA 79  
immunoprecipitation of  $^{35}\text{S}$ -methionine/cysteine labeled  
10 KGLa lysates. In the figure, the faint band at  
~90,000 mw shown on both MECA 79 and rat IgM-control  
antibody immunoprecipitates of KGLa cells is non-  
specific as it can be removed by extensive preclearing  
of lysates with protein-G agarose alone; a similar  
15 faint 90,000 mw band could also be visualized on IgG1  
control immunoprecipitation of  $^{35}\text{SO}_4$ -labeled KGLa  
cells. Of note, the sulfated glycoprotein CD43 (~110  
kDa) was readily immunoprecipitated from KGLa lysates,  
verifying that  $^{35}\text{SO}_4$  was incorporated into newly  
20 synthesized products and that immunoprecipitation  
conditions were appropriate for detection of specific  
radiolabeled KGLa proteins.

#### EXAMPLE 6

25

*Chlorate inhibition of sulfation does not  
block lymphocyte binding to KGLa. As shown herein*

095515-0249-667220-9T85E60

above, the KG1a L-selectin ligand, like the endothelial ligands, requires sialylation for function, as treatment of KG1a with neuraminidase completely abolishes L-selectin-mediated lymphocyte binding. Applicant, therefore, measured the kinetics of recovery of ligand activity following neuraminidase treatment, and chose experimental conditions such to maintain chlorate inhibition of sulfation throughout the period when the KG1a ligand is being re-expressed on the cell membrane. Cell viability following neuraminidase digestion was typically >99% by trypan exclusion.

Initial studies of the kinetics of recovery of binding activity following neuraminidase treatment showed absence of ligand activity for 10-12 hours; thereafter, ligand activity increased steadily with return to baseline levels within 24 hours. Return of binding activity following desialylation was blocked by metabolic inhibition of N-linked glycosylation by tunicamycin and by inhibition of protein synthesis by cycloheximide (Table 3). Of interest, the results of tunicamycin and cycloheximide experiments indicate that the membrane turnover of the L-selectin ligand is relatively slow in resting cells, as culture of KG1a in the presence of these reagents for up to 20 hours did not alter L-selectin ligand activity (Table 3). However, following neuraminidase



treatment of KG1a, both reagents inhibited return of ligand activity without affecting cell viability (trypan blue exclusion >98%), suggesting that re-expression of ligand activity results from de novo synthesis and post-translational processing of ligand and not from transport of pre-formed ligand from intracellular compartments to the membrane surface.

5 The cleavage of sensitive sialic acid epitopes following neuraminidase treatment of KG1a was confirmed by testing for loss of ligand activity in the adherence assay and by measuring the expression of the sialic acid-dependent L60 epitope of CD43 by flow cytometry (Table 3; Figures 7 and 8). L60-negative/ligand activity-negative neuraminidase-treated KG1a were incubated without chlorate (control) or in presence of chlorate for 24 hours to inhibit sulfation throughout the duration of ligand reexpression. Cell viability was >95% in both chlorate-treated and control cell populations (trypan blue exclusion).

10 Twenty-four hours of chlorate treatment (Figure 7C) did not alter the re-expression of ligand activity; indeed, quantitation of adherence assays performed on KG1a 24 hours after neuraminidase treatment revealed that L-selectin ligand activity in chlorate-incubated KG1a cells was slightly higher than that of non-incubated control cells (Table 3). Quantitated lymphocyte adherence to chlorate-incubated KG1a cells

15

20

25

09358146-072199

was 107% +/- 6% (mean +/-SEM) that of neuraminidase-treated KG1a cells not incubated in chlorate. As with control KG1a cells, lymphocyte adherence to sulfate-treated KG1a was L-selectin-specific: it was Ca++-  
5 dependent, was completely inhibited by anti-L-selectin monoclonal antibodies known to block L-selectin adhesive function and by carbohydrate molecules known to bind L-selectin (e.g., fucoidin), and eliminated by phorbol myristate acetate (PMA) treatment of  
10 lymphocytes to induce L-selectin shedding.

Sequential  $^{35}\text{S}$ - $\text{SO}_4$ -pulse radiolabelling studies at eight hour time intervals within the 24 hours of chlorate incubation indicated that sulfation was inhibited throughout the entire incubation period,  
15 as demonstrated by diminished quantities of TCA-precipitable radiolabeled protein counts (chlorate-incubated counts consistently <10% that of non-chlorate treated controls) and of total  $^{35}\text{SO}_4$ -radiolabelled proteins observed by SDS-  
20 PAGE/autoradiography of cell lysates (Figure 9A,B). However, chlorate did not inhibit total protein synthesis as  $^{35}\text{S}$ -methionine/cysteine incorporated TCA-precipitable counts were not significantly different in chlorate and control groups,  $^{35}\text{S}$ - $\text{SO}_4$ -radiolabelling  
25 of CD43 was markedly diminished by chlorate treatment, even in the terminal eight hours of the 24 hour incubation. Similar to the results of recovery of

093546-072199

ligand activity, chlorate treatment did not block re-expression of CD43 to baseline levels (Figure 8C) despite evident sulfation-deficiency (Figure 9).

Monitoring of membrane CD43 recovery was performed

5 using a sialylated epitope of the protein, indicating that sialylation of CD43 (like sialylation of the L-selectin ligand) was not affected by chlorate treatment.

Flow cytometric analysis of CD43 shown in  
10 Figure 8 demonstrated a loss of a sialylation-specific epitope following neuraminidase treatment (Fig. 8B, bottom) relative to control levels (Fig. 8A, top).

Inhibition of sulfation by sodium chlorate is shown in Figure 9 demonstrated by  
15 immunoprecipitation/SDS-PAGE analysis of  $^{35}\text{S}$ -SO<sub>4</sub> incorporation into CD43. Numbers at the left of the figure are molecular weights in kDa.

#### EXAMPLE 7

##### 20 PSGL-1 Analysis

P-selectin glycoprotein ligand-1 (PSGL-1) is a cell surface mucin-like glycoprotein which serves as a ligand for both P- and E-selectin. This protein is expressed on a variety of myeloid cell lines,  
25 including HL60 cells. In order to determine whether the L-selectin ligand activity of KG1a is related to PSGL-1, Applicants performed flow cytometric analysis

09358416-072199  
66T220-91P5560

of PSGL-1 expression on KG1a, blocking studies utilizing both a P-selectin-Ig chimera and an anti-PSGL-1 monoclonal antibody, and enzymatic cleavage and metabolic studies of PSGL-1 and L-selectin ligand activity. These studies reveal that KG1a cells express immunoreactive and functional PSGL-1, however, expression of PSGL-1 is separable from L-selectin ligand activity: (1) blocking of PSGL-1 function using P-selectin-Ig chimera and anti-PSGL-1 monoclonal antibody had no effect on ligand activity of KG1a cells; (2) cleavage with the proteases molarhagin and O-sialoglycoprotease each abrogated PSGL-1 recognition by P-selectin-Ig chimera and L-selectin activity of PSGL-1, but had no effect on L-selectin ligand activity; (3) metabolic inhibition of sulfation by chlorate eliminated P-selectin ligand activity of PSGL-1 but did not alter L-selectin ligand activity; and (4) KG1a sorted for high-level (>95%) and low-level (>5%) expression of PSGL-1 had equivalently high L-selectin ligand activity, and, moreover, HL60 cells were completely devoid of L-selectin ligand activity despite high PSGL-1 expression. These studies indicate that although PSGL-1 is capable of serving as a ligand for E- and P-selectin, it does not itself, display L-selectin ligand activity. The independent relationship between PSGL-1 and L-selectin ligand activity indicates these functions are subserved by

00344.072199

distinct molecules.

Throughout this application various publications or patents are referenced by citation or number. Full citations for the publications  
5 referenced are listed below. The disclosures of these publications and patents in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

10 The invention has been described in an illustrative manner, and it is to be understood that the terminology which has been used is intended to be in the nature of words of description rather than of limitation.

15 Obviously, many modifications and variations of the present invention are possible in light of the above teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as  
20 specifically described.

0935415 022199  
661220 918560

Expression of Surface Molecules on Cell Lines  
utilized in the Lymphocyte Adherence Assay.

RELATIVE EXPRESSION OF MEMBRANE PROTEINS\*

CELL LINE	LYMPHOCYTE ADHERENCE	LINEAGE	CD34	LEA-1	VLA-4	CD44	Sialyl Le <sup>x</sup>	CD43
KG1a	YES	Myeloid	++++	++++	++++	++++	++++	++++
RPMI 8402	NO	Lymphoid	++++	-	++++	+++	-	++++
HL60	NO	Myeloid	-	-	++++	+	-	++++
Nalm 16	NO	Lymphoid	-	-	++++	-	+++	++++
K562	- NO	Erythroid	-	-	-	-	++	+++
Raji	NO	Lymphoid	-	+++	++++	-	-	+

\*Percentage of positive cells as determined by flow cytometric analysis.

- = 0-5% positive  
+ = 6-35% positive  
++ = 36-65% positive  
+++ = 66-95% positive  
++++ = >96% positive

TABLE 1

Table 2. Lymphocyte Adherence to KG1a

<u>LYMPHOCYTE TREATMENT</u>	<u>Mean (SEEM) of Binding (% of Untreated Control)</u>	
EDTA	0.3	(0.3)
Mannose-6-P	5.7	(1.0)
Fucoidin	1.4	(0.4)
PPME	5.4	(0.5)
LAM1-3 mAb	1.9	(0.4)
Anti CD45 mAb	98.7	(6.3)
IgG <sub>1</sub> Control mAb	115.1	(9.0)
PMA	1.1	(0.3)
<u>KG1A TREATMENT:</u>		
Anti CD34 mABs <sup>†</sup>	116.2	(7.7)
Anti CD45 mAb	98.0	(3.6)
IgG <sub>1</sub> Control mAb	101.8	(8.5)
CD34-Positive Sort	102.8	(3.5)
CD34-Negative Sort	104.1	(4.2)
Neuraminidase	3.1	(0.7)
Neuraminidase Buffer Control	100.5	(6.7)
O-Sialoglycoprotein Endopeptidase	98.4	(2.3)
Bromelain	3.8	(0.4)
Chymotrypsin	6.7	(0.7)
Chymotrypsin, PMSF, Chymostatin	94.0	(3.8)
Combination of HPCA-1, HPCA-2, 12.8 and QBEND10 mAbs.		

09353115.072109

TABLE 3

## Lymphocyte Adherence to KG1a

KG1a Treatment	Mean (SEM) of Binding (% of untreated control)**
Neuraminidase (t=0, no culture)	0.02 (0.1)
Neuraminidase Buffer Control (no culture)	98.5 (7.5)
Neuraminidase, 24 hr culture	101.3 (6.5)
Neuraminidase, 24 hr culture with Chlorate (10 mM)	107.1 (6.2)
Neuraminidase, 20 hr culture with Tunicamycin (15 ug/ml)	0.4 (0.1)
Neuraminidase, 20 hr culture with Tunicamycin Buffer alone	105.2 (7.5)
20 hr culture with Tunicamycin (15 ug/ml) (no neuraminidase pretreatment)	106.2 (7.2)
Neuraminidase, 20 hr culture with Cycloheximide (1.25 ug/ml)	0.5 (0.2)
20 hr culture with Cycloheximide (1.25 ug/ml) (no neuraminidase pretreatment)	91.8 (5.8)

\*Following neuraminidase treatment, all cells were washed prior to culturing as indicated. Experimental details are described in text.

\*\*Number of lymphocytes adherent to confluent area of KG1a were counted by light microscopy using an ocular grid under 250X magnification (quantified a minimum of 2 fields/slide, 2 slides/experiment, 3 separate experiments). Results are presented as percent binding compared with corresponding untreated control KG1a cyto-spin preparations.

0152.00344



REFERENCES

- Abdullah et al., "A neutral glycoprotease of *Pasteurella haemolytica* A1 specifically cleaves O-sialoglycoproteins" Infect Immun., 60:56, (1992).
- Aruffo et al., "CD62/P-selectin recognition of myeloid and tumor cell sulfatides" Cell, 67:35-44 (1991).
- Baeuerle and Huttner, "Chlorate-a potent inhibitor of protein sulfation in intact cells" Biochemical and Biophysical Res. Comm., 141:870-877 (1986).
- Baumhueter et al., "Binding of L-selectin to the vascular sialomucin CD34" Science 262:436, (1993).
- Berg et al., "L-selectin mediated lymphocyte rolling on MAD-CAM-1" Nature, 366:695-698 (1993).
- Beschorner et al., "Localization of hematopoietic progenitor cells in tissue with the anti-My-10 monoclonal antibody" Am J Pathol 119:1, (1985).
- Bevilacqua and Nelson, "Selectins" J Clin Invest., 91:379, (1993).
- Bollman et al., "Techniques for the collection of lymph from the liver, small intestine, or thoracic duct of the rat" J Lab Clin Med., 33:1349, (1948).
- Carlos and Harlan, "Leukocyte-Endothelial adhesion molecules" Blood, 84(7):2068, (1994).
- Chin et al., "Lymphocyte-homing receptors and preferential migration pathways" Proc Soc Exp Biol Med., 196:374, (1991).
- Civin et al., "Antigenic analysis of hematopoiesis III. A hematopoietic progenitor cell surface antigen defined by a monoclonal antibody raised against KG-1a cells" J Immunol., 133:157, (1984).

095816-07199

Cowing and Chapdelaine, "T cells discriminate between Ia antigens expressed on allogeneic accessory cells and B cells: a potential function for carbohydrate side chains on Ia molecules" Proc Natl Acad Sci USA, 80:6000, (1983).

Cullen et al., "Differential glycosylation of murine B cell and spleen adherent cell Ia antigens" J Immunol., 127:1478, (1981).

Cytospin®3 Cell Preparation System Operator Guide, Shandon Scientific Limited.

Dercksen et al., "Expression of adhesion molecules on CD34+ cells: CD34+ L-selectin+ cells predict a rapid platelet recovery after peripheral blood stem cell transplantation" Blood, 85:3313-3319 (1995).

Dowbenko et al., "Structure and chromosomal localization of the murine gene encoding GlyCAM-1" J. Clin. Invest., 92:952-960 (1993).

Gallatin, et al., "A cell-surface molecule involved in organ-specific homing of lymphocytes" Nature, 304:30-34 (1983).

Gatenby et al., "Dissection of immunoregulatory subpopulations of T lymphocytes within the helper and suppressor sublineages in man" J. Immunol., 129:1997, (1982).

Gowans and Knight, "The route of re-circulation of lymphocytes in the rat" Proc Roy Soc Lond B., 159:257, (1964).

Green et al., "Further studies of the binding specificity of the leukocyte adhesion molecule, L-selectin, towards sulphated oligosaccharides-suggestion of a link between the selectin- and the integrin-mediated lymphocyte adhesion systems" Glycobiology, 5:29-38 (1995).

Green et al., "High affinity binding of the leukocyte adhesion molecule L-selectin to 3'-sulphated-Le<sup>x</sup> and -Le<sup>x</sup> oligosaccharides and the predominance of sulphate in this interaction demonstrated by binding studies with a series of lipid-linked oligosaccharides" Biochem. Biophys. Res. Comm., 188:244-251 (1992).

035815-074560

Griffin et al., "Granulocyte-macrophage colony-stimulating factor and other cytokines regulate surface expression of the leukocyte adhesion molecule-1 on human neutrophils, monocytes, and their precursors" J Immunol., 145:576, (1990).

Hemmerich, et al., "Sulfation-dependent recognition of high endothelial venules (HEV)-ligands by L-selectin and MECA79, an adhesion blocking monoclonal antibody" J. Exp. Med., 180:2210-2226 (1994).

Hemmerich and Rosen, "6'-sulfated sialyl lewis x is a major capping group of GlyCAM-1" Biochemistry, 33:4830-4835 (1994b).

Huff et al., "Modulation of the biologic activities of IgE-binding factors. III. switching of a T cell hybrid clone from the formation of IgE-suppressive factor to the formation of IgE-potentiating factor" J Immunol., 131:1090, (1983).

Imai et al., "Sulphation requirement for GlyCAM-1, an endothelial ligand for L-selectin" Nature 361:555, (1993).

Imai et al., "Identification of a carbohydrate-based endothelial ligand for a lymphocyte homing receptor" J Cell Biol., 113:1213, (1991).

Kobayashi et al., "Expression of adhesion molecules on human hematopoietic progenitor cells at different maturational stages" Stem Cells, 12:316, (1994).

Koeffler et al., "An undifferentiated variant derived from the human acute myelogenous leukemia cell line (KG-1)" Blood 56:265, (1980).

Koenig et al., Blood, 84:599 (1994).

Krause et al., "Two forms of CD34 protein are expressed in human KMT2 and KG1a cells" Blood, 82:110a, (1993) (abstr).

Lasky et al., "An endothelial ligand for L-selectin is a novel mucin-like molecule" Cell, 69:927, (1992).

00344-0152

Lewinsohn et al., "Leukocyte-endothelial cell recognition: evidence of a common molecular mechanism shared by neutrophils, lymphocytes, and other leukocytes" J Immunol., 138:4313, (1987).

Majdic et al., "Signaling and induction of enhanced cytoadhesiveness via the hematopoietic progenitor cell surface molecule CD34" Blood, 83:1226, (1994).

Marchesi and Gowans, "The migration of lymphocytes through the endothelium of venules in lymph nodes: an electron microscope study" Proc Roy Soc Lond B, 159:283, (1964).

Niehurs et al., Chemico-Bio. Interactions, 92:257-271 (1992).

Ord et al., "Structure of the gene encoding the human leukocyte adhesion molecule-1 (TQ1, Leu-8) of lymphocytes and neutrophils" J Biol Chem., 265:7760, (1990).

Oxley and Sackstein, "L-selectin-CD34 Interactions mediate functional binding of lymphocytes to hematopoietic progenitor cells" Clin Res. Abs. 42(2):235A, April (1994).

Oxley and Sackstein, "Detection of an L-selectin ligand on a hematopoietic progenitor cell line" Blood, 84:3299-3306 (1994b)

Pouyani and Seed, "PSGL-1 recognition of P-selectin is controlled by a tyrosine sulfation consensus at the PSGL-1 amino terminus" Cell, 83:333-343 (1995).

Puri et al., "Sialomucin CD34 is the major L-selectin ligand in human tonsil high endothelial venules" J. Cell Bio., 131:261-270 (1995).

Ramakrishnan, "Current status of antibody-toxin conjugates for tumor therapy" Targeted Diagn Ther., 3:189, (1990).

Rasmussen et al., "Lymphocyte recognition of lymph node high endothelium. VII. Cell surface proteins involved in adhesion defined by monoclonal anti-HEBFLN (A.11) antibody" J. Immunol., 135:19-24 (1985).

0152.00344

Reinherz et al., "Heterogeneity of human T4<sup>+</sup> inducer T cells defined by a monoclonal antibody that delineates two functional subpopulations" J Immunol., 128:463, 1982.

Rosen et al., "Involvement of sialic acid on endothelial cells in organ-specific lymphocyte recirculation" Science, 228:1005, (1985).

Rosen, "Cell surface lectins in the immune system" Sem Imm., 5:237, (1993).

Rosen, "Localization of ligands for L-selectin in mouse peripheral lymph node high endothelial cells by colloidal gold conjugates" Blood, 84:3766-3775 (1994).

Sackstein, et al., "Evidence of post-transcriptional regulation of L-selectin gene expression in rat lymphoid cells" Immunol., 85:198-204 (1995).

Sackstein, "Physiological migration of lymphocytes to lymph nodes following bone marrow transplantation: Role in immune recovery" Seminars in Oncology 20(5), Suppl 6:34, (1993).

Sackstein et al., "Lymphocyte adhesion to psoriatic dermal endothelium is mediated by a tissue-specific receptor/ligand interaction" J Invest Dermatol., 91:423, (1988).

Sako et al., "A sulfated peptide segment at the amino terminus of PSGL-1 is critical for P-selectin binding" Cell, 83:323-331 (1995).

Selden, "Transfection using DEAE-dextran", Ausubel FM, Brent R, Kingston RE, Moor DD, Seidman JG, Smith JA, Struhl K (eds) in Current Protocols in Molecular Biology, Vol 2. New York NY, John Wiley & Sons, p. 9.2.1. (1992).

Sharon and Lis, "Carbohydrates in cell recognition" Sci. Amer., 82-89, Jan. (1993).

Shaw et al., "Two antigen-independent adhesion pathways used by human cytotoxic T-cell clones" Nature, 323:262, (1986).

033416-034109

Shimazu and Shaw, "Mucins in the mainstream" Nature 366:630-631 (1993).

Spertini et al., "Leukocyte adhesion molecule-1 (LAM-1, L-selectin) interacts with an inducible endothelial cell ligand to support leukocyte adhesion" J Immunol., 147:222565, (1991).

Spertini et al., "Function and evolutionary conservation of distinct epitopes on the leukocyte adhesion molecule-1 (TQ1, Leu-i) that regulate leukocyte migration" J Immunol., 147:942 (1991).

Springer et al., "The lymphocyte function-associated LFA-1, CD2, and LFA-3 molecules: Cell adhesion receptors of the immune system" Ann Rev Immunol., 5:223-252 (1987).

Stamper and Woodruff, "Lymphocyte homing into lymph nodes: in vitro demonstration of the selective affinity of recirculating lymphocytes for high-endothelial venules" J Exp Med., 144:828-833 (1976).

Stoolman et al., "Phosphomannosyl receptors may participate in the adhesive interaction between lymphocytes and high endothelial venules" J Cell Biol., 99:1111535 (1984).

Stoolman and Rosen, "Possible role for cell-surface carbohydrate-binding molecules in lymphocyte recirculation" J Cell Biol., 96:722 (1983).

Streeter et al., "A tissue-specific endothelial cell molecule involved in lymphocyte homing" Nature, 331:41 (1988).

Sutherland et al., "Differential sensitivity of CD34 epitopes to cleavage by *Pasteurella haemolytica* glycoprotease: Implications for purification of CD34-positive progenitor cells" Exp Hematol., 20:590 (1992).

Suzuki et al., "Sulfated glycolipids are ligands for a lymphocyte homing receptor, L-selectin (LECAM-1), binding epitope in sulfated sugared chain" Biochem. Biophys. Res. Comm., 190:426-434 (1993).

Tedder et al., "Expression of the human leukocyte adhesion molecule, LAM1, identity with the TQ1 and Leu-8 differentiation antigens" J Immunol., 144:532-540 (1990).

09355816-02199

Terstappen et al., "Induction of human hematopoietic stem cell outgrowth by L-selectin ligation" Blood 82:109a (1993) (abstr)

Terstappen et al., "Flow cytometric assessment of human T-cell differentiation in thymus and bone marrow" Blood 79:666 (1992).

True et al., "Requirement for sialic acid on the endothelial ligand of a lymphocyte homing receptor" J Cell Biol., 111:2757 (1990).

Watson et al., "A homing receptor-IgG chimera as a probe for adhesive ligands of lymph node high endothelial venules" J Cell Biol., 110:2221 (1990).

Wilson and Rider, "Evidence that Leukosialin, CD43, is intensely sulfated in the murine T lymphoma line RDM-4" J. Immunol., 148:1777-1783 (1992).

Woodruff et al., "An in vitro model of lymphocyte homing II. Membrane and cytoplasmic events involved in lymphocyte adherence to specialized high-endothelial venules of lymph nodes" J Immunol., 119:1603 (1977).

Woodruff et al., "Specific cell-adhesion mechanisms determining migration pathways of recirculating lymphocytes" Ann Rev Immunol., 5:201-222 (1987).

Yamashita et al., "Organ-specific difference in the sugar chains of gamma-glutamyltranspeptidase. Arch Biochem Biophys., 225:993 (1983).

Yamashita et al., "The structures of the carbohydrate moieties of mouse kidney gamma-glutamyltranspeptidase: occurrence of X-antigenic determinants and bisecting N-acetylglucosamine residues" Arch Biochem Biophys., 240:573 (1985).

Yednock et al., "Phosphomannosyl-derived beads detect a receptor involved in lymphocyte homing" J. Cell Biol., 104:713-723 (1987).

093546 07219  
66T20 9T95660